arXiv:1509.04519v2 [math.CO] 2 Feb 2017

On enumeration of a class of maps on Klein bottle

Dipendu Maity

School of Mathematics
Harish-Chandra Research Institute
Chhatnag Road, Jhunsi Allahabad-211019, India.
dipendumaity@hri.res.in

Ashish Kumar Upadhyay

Department of Mathematics
Indian Institute of Technology Patna
Patliputra Colony, Patna 800 013, India.
upadhyay@iitp.ac.in

February 3, 2017

Abstract

There are eleven types of semi-equivelar maps on the torus and the Klein bottle [5]. Three of these are eqivelar maps and the remaining are semi-equivelar maps. We classify all eleven types of semi-equivelar maps on the Klein bottle.

AMS classification : 52C20, 52B70, 51M20, 57M60

Keywords : Polyhedral maps on Klein bottle, Vertex-transitive maps, Semi-Equivelar Maps

1 Introduction and definitions

The following definitions one can see in [3, 5, 7] and graph related terms in Bondy and Murthy [3]. These are being reproduced here for the sake of completeness. A *map* M is an embedding of a finite graph G on a surface S such that the closure of components of $S \setminus G$, called the faces of M, are closed 2-cells, that is, each homeomorphic to 2-disk. By a map we mean a polyhedral map on the torus. A *n-cycle* C_n is a finite connected 2-regular graph with n vertices, and the *face sequence* of a vertex v in a map K is a finite sequence $(a^p, b^q, \ldots, m^r)$ of powers of positive integers $a, b, \ldots, m \geq 3$ and $p, q, \ldots, r \geq 1$ in *cyclic* order such that through the vertex v, p number of C_a (C_a denote the a-cycle), q number of C_b, $\ldots$, r number of C_m are incident. In this case, we say that K is *semi-equivelar* map of type $\{a^p, b^q, \ldots, m^r\}$ [5, 7]. A map of type $\{p^q\}$ is a semi-equivelar map of type $\{p^q\}$. We say two maps of fixed type on the Klein bottle are *isomorphic* if there exists a *homeomorphism* of the Klein bottle which sends vertices to vertices, edges to edges, faces to faces and preserves incidents. More precisely, if we consider two polyhedral complexes K_1 and K_2 then an isomorphism to be a map $f : K_1 \to K_2$ such that $f|_{V(K_1)} : V(K_1) \to V(K_2)$ is a bijection and $f(\sigma)$ is a cell in K_2 if and only if σ is cell in K_1.

There are eleven types of semi-equivelar maps on the torus and the Klein bottle, [5]. These types are $\{3^6\}, \{4^4\}, \{6^3\}, \{3^3, 4^2\}, \{3^2, 4, 3, 4\}, \{3, 6, 3, 6\}, \{3^4, 6\}, \{4, 8^2\}, \{3, 12^2\}, \{4, 6, 12\}, \{3, 4, 6, 4\}$. In 1973, Altshuler [2] has shown the construction and enumeration of maps of types $\{3^6\}$ and $\{6^3\}$ on the torus. He has studied the maps of type $\{3^6\}$ on the torus with n vertices and has given a characterization of these maps by introducing some arithmetic conditions. In 1986, Kurth [6] has given enumeration of all equivelar maps of types $\{3^6\}, \{4^4\}$ and $\{6^3\}$ on the torus using quotients of planar tessellations by squares or equilateral triangles, respectively, under the action of a group of isometries generated by two translations. In 1983, Negami [10] has given uniqueness and faithfulness of embedding of 6-regular toroidal graphs and then classified the maps of type $\{3^6\}$ by classifying the 6-regular graphs on the torus. Similarly in 2008, Brehm and Kühnel [4] have presented a classification of equivelar maps of types $\{3^6\}, \{4^4\}, \{6^3\}$ on the torus using their different isotropy groups. In [7], Maity and Upadhyay have given a construction to enumerate all semi-equivelar maps of types $\{3^3, 4^2\}, \{3^2, 4, 3, 4\}, \{3, 6, 3, 6\}, \{3^4, 6\}, \{4, 8^2\}, \{3, 12^2\}, \{4, 6, 12\}, \{3, 4, 6, 4\}$ on the torus for arbitrary number of vertices. By the Euler characteristic equation, it is easy to see that there is no bound on number of vertices of the toroidal maps. Therefore, the above articles have presented computational steps or formulas to calculate the non-isomorphic maps on the torus. Thus, by [2, 4, 6, 7, 10, 11], we know

Proposition 1.1. *Let n denote the number of vertices of maps of type $\{3^6\}, \{4^4\}, \{6^3\}, \{3^3, 4^2\}, \{3^2, 4, 3, 4\}, \{3, 6, 3, 6\}, \{3^4, 6\}, \{4, 8^2\}, \{3, 12^2\}, \{4, 6, 12\}$ or $\{3, 4, 6, 4\}$ on the torus. Then we can classify maps on n vertices upto isomporphism.*

This paper is devoted to a study of all semi-equivelar maps on the Klein bottle with n vertices. We devise a way (computational steps) to enumerate and characterise all semi-equivelar maps of types $\{3^6\}, \{4^4\}, \{6^3\}, \{3^3, 4^2\}, \{3^2, 4, 3, 4\}, \{3, 6, 3, 6\}, \{3^4, 6\}, \{4, 8^2\}, \{3, 12^2\}, \{4, 6, 12\}, \{3, 4, 6, 4\}$ on the Klein bottle. In next theorem we give a formula on n number of vertices which enumerate number of non-isomorphic maps on the Klein bottle.

Theorem 1.1. *Let $X \in \{\{3^6\}, \{4^4\}, \{6^3\}, \{3^3, 4^2\}, \{3^2, 4, 3, 4\}, \{3, 6, 3, 6\}, \{3^4, 6\}, \{4, 8^2\}, \{3, 12^2\}, \{4, 6, 12\}, \{3, 4, 6, 4\}\}$. Let $i(n)_X$ denote the number of non-isomorphic semi-equivelar maps with n vertices of type X on the Klein bottle. Then,* (1) $i(n)_{\{3^6\}} = \sum_{i=1,2} i \times |\{(m, n) \mid m \geq 3, n \geq 3m,\ gcd(n, 2m) = i \times m\}| + |\{(m, n) \mid m \geq 2, m \mid n, n \geq 5m,\ gcd(n, 2m) = m\}|$, (2) $i(n)_{\{4^4\}} = \sum_{i=1,2} i \times |\{(m, n) \mid m \geq 3, n \geq 3m, gcd(n, 2m) = i \times m\}|$, (3) $i(n)_{\{6^3\}} = i(\frac{n}{2})_{\{3^6\}}$, (4) $i(n)_{\{3^3,4^2\}} = \sum_{i=1,2} i \times |\{(m, n) \mid m \geq 4, 2 \mid m, n \geq 3m, gcd(n, 2m) = m\}| + |\{(m, n) \mid m \geq 5, n = 2m\}| + |\{(m, n) \mid m \geq 4, n \geq 4m, 2m \mid n\}|$, (5) $i(n)_{\{3^2,4,3,4\}} = |\{(m, n) \mid m \geq 3, 2 \nmid m, 2m \mid n, n \geq 12\}|$, (6) $i(n)_{\{4,8^2\}} = |\{(l, m, n) \mid m \geq 3, 2 \mid m, n \geq 8m, 4m \mid n, 0 \leq l \leq (\frac{n}{4m} - 1)\}| + |\{(l, m, n) \mid m \geq 3, 2 \nmid m, n \geq 8m, 4m \mid n, 0 \leq l \leq (\frac{n}{4m} - 1)\}|$, (7) $i(n)_{\{3,6,3,6\}} = |\{(l, m, n) \mid m \geq 3, 3m \mid n, n \geq 9m, 0 \leq l \leq (\frac{n}{3m} - 1)\}| + |\{(m, n) \mid m \geq 2, 6m + 2 \mid n, n \geq 6(3m + 1)\}| + |\{(m, n) \mid m \geq 1, 2(m + 2) \mid n, n \geq 10(m + 2)\}| + |\{(m, n) \mid m \geq 1, 4m + 5 \mid n, n \geq 12(4m + 5)\}|$, (8) $i(n)_{\{3,12^2\}} = |\{(l, m, n) \mid m \geq 3, 3m \mid n, n \geq 9m, 0 \leq l \leq (\frac{n}{6m} - 1)\}| + |\{(m, n) \mid m \geq 2, 4(3m + 1) \mid n, n \geq 12(3m + 1)\}| + |\{(m, n) \mid m \geq 1, 4(m + 2) \mid n, n \geq 20(m + 2)\}| + |\{(m, n) \mid m \geq 1, 4m + 5 \mid n, n \geq 24(4m + 5)\}|$, (9) $i(n)_{\{3^4,6\}} = 0$, (10) $i(n)_{\{4,6,12\}} = \{(l, m, n) \mid m \geq 4, 2 \mid m, 6m \mid n, n \geq 12m, 0 \leq l \leq (\frac{n}{6m} - 1)\}| + |\{(m, n) \mid m \geq 2, 2 \mid m, 12m \mid n, n \geq 24m\}|$, (11) $i(n)_{\{3,4,6,4\}} = |\{(l, m, n) \mid m \geq 4, 2 \mid m, 3m \mid n, n \geq 6m, 0 \leq l \leq (\frac{n}{3m} - 1)\}| + |\{(m, n) \mid m \geq 2, 6m \mid (n - 3m), n \geq 9m\}| + |\{(m, n) \mid m \geq 2, 6m \mid n, n \geq 12m\}|$.

We introduced some specific types of cylindrical polyhedral representations which are bounded by identical cycles with different orientations or by mobius strips that are consists of polygonal faces of semi-equivelar maps on the Klein bottle. By these polyhedral representations, we give a complete characterizations of all semi-equivelar maps on the Klein bottle. So, we have

Theorem 1.2. *Let M be a map of type X with n vertices on the Klein bottle. Then, the map M is isomorphic to a polyhedral representation $K(r, s, 0), K(r, s, 1)$ or $K(r, s)$ (these polyhedral representations are defined later in the subsequent sections) for some $r \mid n$ and $s = f(n, r)$.*

Remark 1.1. In Theorem 1.2, the $K(r, s, 0), K(r, s, 1)$ are called planar polyhedral representations which are bounded by identical cycles with different orientations and $K(r, s)$ is called polyhedral representations which is bounded by mobius strips. We have computed all eleven types of semi-equivelar maps with on few number of vertices in [9] using the idea of the proof of Theorem 1.2.

The proofs of all above theorems are presented in Section 3. We study above three polyhedral representations of all eleven types of semi-equivelar maps in the subsequent sections. We use the results of the following sub-sections of Section 2 in Section 3 and prove our main results.

2 Semi-equivelar Maps

2.1 Maps of type $\{3^6\}$: Let M be a map of type $\{3^6\}$ on the Klein bottle. A path $P(\dots, u_{i-1}, u_i, u_{i+1}, \dots)$ in edge graph of M is of type A if $lk(u_i) = C(a, b, u_{i+1}, c, d, u_{i-1})$ implies $lk(u_{i-1}) = C(g, a, u_i, d, u, u_{i-2})$ and $lk(u_{i+1}) = C(b, e, u_{i+2}, f, c, u_i)$ for all i. A walk $Walk(u_1, e_{1,2}, u_2, \dots, u_i, e_{x-1,x}, u_x)$ in edge graph of M is a sequence of vertices and edges such that $e_{t-1,t} = u_{t-1}u_t, 1 \leq t \leq x$. We denote it by $Walk(u_1, u_2 \dots, u_x)$.

Claim : If $W(w_1, \dots, w_r)$ is a maximal walk of type A in $EG(M)$ (edge graph of M) then $W(w_1, \dots, w_r)$ is a closed walk of type A.

By the defintion of A, let $lk(w_{r-1}) = C(a, b, w_r, c, d, w_{r-1})$, it implies that $lk(w_r) = C(b, e, w_{r+1}, f, c, w_r)$. If $w_r = w_1$ then $C(w_1, w_2, \dots, w_r)$ is a closed walk of type A. If $w_r \neq w_1$, it implies that $w_r = w_i$ for some $2 \leq i \leq r$. Hence, we get a closed walk $R = Walk(w_i, w_{i+1}, \dots, w_r)$ and which is not of type A at w_i. This implies that the number of triangles which are incident at w_i is one or two on one side of R. If the number of incident triangles is one then we define an another closed walk as R of lesser length using the incident faces of the walk W. (The similar description of this construction is given details for a cycle in *Theorem 1* [1] (also see it in the example in *Fig. 2* [1]).) Similarly, if the number triangles which are incident at w_i is two then one can see that we get a cycle of same type (as in *Fig. 3* [1]). Thus, we get a closed walk as R of lesser length using the triangles which are incident with R. By induction on length, it is not possible to construct walk as R at each step as length is decreasing. Therefore, $w_r = w_1$ and $W(w_1, \dots, w_r)$ is a closed walk of type A. This proves the claim.

Claim : A closed walk $W(w_1, \dots, w_r)$ of type A is non-contractible.

If the walk W bounds a 2-disk then we consider triangles which are incident with W in the 2-disk. There are three triangles which are incident at each vertex of W in the 2-disk. Let $w_{i-1}w_iv_i, w_{i-1}v_iv_{i+1}, w_iv_{i+1}w_i$ denote three triangles incident at w_i for all i. These incident triangles define a walk $W' = Walk(v_1, \dots, v_r)$ of type A and has a length r as W. Similarly, we consider walk W' and repeat as above. Hence, we get an infinitely many sequence of walks namely $W, W', W'', \ldots$ where each walk has length r. But, the 2-disk consists of finite number of triangles. Therefore, the walk W does not bound any 2-disk. Which is a contradiction. So, W is non-contractible. This proves the claim. We show in the next lemma that the map M contains a cycle of type A.

Lemma 2.1. *The map M contains a cycle of type A.*

Proof. Let W be a closed walk of type A in edge graph of M and S_W denote a set of faces which are incident with walk W. Observe that by considering incident faces of W, the boundary of the geometric carrier S_W consists of two closed walks. It follows that $\partial S_W = \{W_2, W_s\}$ where W_2 and W_s are two closed walks of type A and have same length. Here we say that the walks W_2, W_s are homologous to W. Since W is a walk, let $w \in V(W)$ repeat twice in W. Observe that by the definition of A, there is no repetition of edges in W. Since $w \in V(W)$ repeats twice in W, it follows that there are two sub-paths $P_1 = P(w_{l-1}, w, w_{l+1})$ and $P_2 = P(w_{k-1}, w, w_{k+1})$ of W where $w_{l-1}, w_{l+1}, w_{k-1}, w_{k+1} \in lk(w)$. Let $v \in lk(w)$. Let $W_1 (= W), W_2, \ldots, W_s$ be a sequence of walks which are homologous to W in M.

Claim : The vertex v repeats twice in $W_1 \cup W_2 \cup \cdots \cup W_s$

If $P_1 \subset W$ then $v \in W_i$ for some i and $W_i \in \partial S_W$. Similarly, if $P_2 \subset W$ then $v \in W_j$ for some j and $W_j \in \partial S_W$. So, the vertex v repeats twice in $W_i \cup W_j$. Therefore, every vertex of M repeats twice in $W_1 \cup W_2 \cup \cdots \cup W_s$ since $u \in W_1 \cup W_2 \cup \cdots \cup W_s \ \forall \ u \in V(M)$. This proves the Claim.

Similarly as in the Claim, if a vertex repeats thrice in W then every vertex of M repeats thrice in $W_1 \cup W_2 \cup \cdots \cup W_s$. Since W is non-contractible, it implies that the W represents a generator of the fundamental group of M. The fundamental group of M contains exactly two generator up to homologous. Let W', W'' be two non-homologous closed walks of type A through a vertex in M. Here, if the walk W' contains a vertex, say w which repeats twice or thrice in W' then we have following cases. If the vertex w repeats thrice then by the above Claim, every vertex of M repeats thrice in the union of homologous walks of W'. This implies that the union of walks contains all the edges of M. Which is not possible since W', W'' are non-homologus and which implies that $E(W') \cap E(W'') = \emptyset$. So, walk W' contains the vertex w twice. Let $L_1, \ldots, L_l$ denote sequence of homologous walks of W' in M. Since W' contains the vertex w twice, it implies by Claim that $L_1 \cup \cdots \cup L_l$ contains each vertex twice of M. Again, $E(W') \cap E(W'') = \emptyset$ implies that $E(W'') \cap E(L_i) = \emptyset$ for all $1 \leq i \leq l$. Hence, walk W'' contains all distinct vertices since degree of each vertex of M is six. So, W'' is a cycle. Therefore, map M contains a cycle of type A. $\square$

For an example, the $W(v_1, w_1, x_1, u_1, v_2, w_3, x_4, u_5, u_4, w_4, x_4, u_4, v_6, w_7, x_1, u_2, v_7, w_7, x_7, u_7, v_3, w_4, x_5, u_6, v_3, w_3, x_3, u_3, v_7, w_1, x_2, u_3, v_6, w_6, x_6, u_6, v_4, w_5, x_6, u_7, v_2, w_2, x_2, u_2, v_1, w_2, x_3, u_4, v_5, w_5, x_5, u_5, v_5, w_6, x_7, u_1, v_1)$ is a closed walk of type A and $C(x_1, \ldots, x_7)$ is a cycle of type A in Figure 1. In W each vertex repeats exactly twice.

Let C be a cycle of type A in M and $u \in V(C)$. A set $S \subset F(M)$ is of faces such that $V(\triangle) \cap V(C) \neq \emptyset$ for all $\triangle \in S$. Then, we have

Lemma 2.2. *The geometric carrier $|S|$ is a cylinder or an union of 'a Möbius strip and a cylinder'.*

Proof. Let SQ be a sequence of faces which are incident with the cycle C. Let $u_1 \in V(C)$ and $lk(u_1) = C(v_r, v_1, u_2, w_2, w_1, u_r)$. We cut $|SQ|$ along a path $P(v_1, u_1, w_1)$ of type A. Then we have the following cases.

Case 1 : We assume that each face in SQ appears exactly once. Here, we have the following sub-cases. If the path $P(v_1, u_1, w_1)$ identifies with $P(v_1, u_1, w_1)$ with out any twist then $|SQ|$ is cylinder (see an example in Figure 2). In this acse, we denote it by $S(C, M)$. If $P(v_1, u_1, w_1)$ identifies with $P(v_1, u_1, w_1)$ with a twist then the cycle C is not of type A at the vertex u_1. (For an example in Figure 3, $C(w_1, \dots, w_r)$ is a cycle and not of type A at w_1.) Therefore, $|S| = |SQ| = S(C, M)$ is a cylinder if the faces are distinct in SQ.

Case 2 : We assume that not all faces are distinct in SQ. Let SQ' denote an another sequence of faces which are incident with C and lie on one side of C. Here, the length of C is either odd or even.

Let $C = C(w_1, w_2, \dots, w_{2r+1})$ and $SQ' = \{\triangle_1, \triangle_2, \dots, \triangle_{4r+2}\}$ denote the sequence of faces in order. If $\triangle_1, \triangle_2, \dots, \triangle_{4r+2}$ are all distinct, i.e., $\triangle_i \neq \triangle_j$ for all $i \neq j$ then $|SQ'|$ forms a cylinder with disjoint boundary cycles. If $\triangle_i = \triangle_j$ in SQ' for some $i < j$ then $SQ'' := \{\triangle_1, \triangle_2, \dots, \triangle_t\} \subset SQ'$ denotes a subsequence where $\{\triangle_1, \triangle_2, \dots, \triangle_t\} = \{\triangle_1, \triangle_2, \dots, \triangle_{4r+2}\}$ as a set, and $\triangle_1$ and $\triangle_t$ appear once in $\{\triangle_1, \triangle_2, \dots, \triangle_t\}$.

Claim : The triangles $\triangle_1, \triangle_2, \dots, \triangle_t$ appear once in SQ''.

Let $\{\triangle_l, \dots, \triangle_s\}$ denote a largest sub-sequence in SQ'' where each $\triangle_l, \dots, \triangle_s$ appears more than once in SQ''. This implies that the faces $\triangle_{l-1}$ and $\triangle_{s+1}$ are distinct and appear once in SQ''. Let $\triangle_s = abc$ and $\triangle_s \cap C = ab$. Then, $\triangle_s \cap \triangle_{s+1} = bc$. Since $\triangle_s$ repeats in SQ'', it implies that there are $\triangle_{m-1}$ and $\triangle_{m+1}$ in SQ'' such that $\triangle_m = \triangle_s$ $(m < s)$. That is, $\triangle_s$ lies between $\triangle_{m-1}$ and $\triangle_{m+1}$ in SQ''. Hence, $\triangle_s \cap \triangle_{m-1}$ and $\triangle_s \cap \triangle_{m+1}$ are each an edge. Since $\triangle_{s+1}$ appears first time in SQ'' after $\triangle_s$, it follows that either $\triangle_{s-1} \neq \triangle_{m-1}$ or $\triangle_{s-1} \neq \triangle_{m+1}$. If $\triangle_{s-1} \neq \triangle_{m+1}$ then $\triangle_{s+1} \neq \triangle_{s-1} \neq \triangle_{m+1}$. Now, by definition of type A, one edge, say e of $\triangle_s$ must belongs to C or C'. Hence, the triangle $\triangle_s$ must have at least four distinct edges $\triangle_{s+1} \cap \triangle_s, \triangle_{s-1} \cap \triangle_s, \triangle_{m+1} \cap \triangle_s, e$. This is a contradiction as triangle consists three edges. Similarly, let $\triangle_s \cap C = a$. Then by the above similar argument, $\triangle_s$ does not have sufficient number of free edges for identification later in SQ''. Thus, the triangles $\triangle_1, \triangle_2, \dots, \triangle_t$ appear once in SQ''. This proves the Claim.

Let $\triangle_i = ABC \in SQ'$. Now $\triangle_i \cap \triangle_{i-1}$ and $\triangle_i \cap \triangle_{i+1}$ are each an edge. Let $\triangle_i \cap \triangle_{i-1} = AB$ and $\triangle_i \cap \triangle_{i+1} = AC$. Hence, $\triangle_i \cap C$ is either an edge BC or a vertex. Let $\triangle_i \cap C = BC$. Then, $\triangle_i \cap C'$ is a vertex. Hence, the common edge BC and the common vertex can not repeat twice in both C, C' since C, C' are cycles. But $\triangle_i$ repeats twice in SQ' only if $C = C'$. Since $\triangle_i$ is repeating, it implies that it is repeating twice. Hence, each triangle appears twice in the sequence SQ'. Thus, the number of distinct triangle in SQ' is $2r + 1$. That is, $t = 2r+1$. So, the sequence $\triangle_1, \triangle_2, \dots, \triangle_{2r+1}$ appears twice in the sequence $\triangle_1, \triangle_2, \dots, \triangle_{4r+2}$, i.e., $\{\triangle_1, \triangle_2, \dots, \triangle_{4r+2}\} = \{\triangle_1, \triangle_2, \dots, \triangle_{2r+1}, \triangle_1, \triangle_2, \dots, \triangle_{2r+1}\}$. In this case, we call that $|SQ'|$ is a Möbius strip. (See an example in Figure 4.)

When length$(C) = 2r$ then the number of triangles incident with SQ' is even. In this case, we consider link of w_{r+1} which is end vertex of the path $P(w_1, \dots, w_{r+1}) \subset C(w_, w_2, \dots, w_{2r})$. Observe that the number of triangles which are incident with C at w_{r+1} is two in SQ'. This contradicts with the type A.

Let S_1 denote a set of incident triangles which are lie on one side and S_2 denote a set of triangles which are incident with C and lie on the other side of C. If $S_1 \cap S_2 \neq \emptyset$ then, let $\triangle \in S_1 \cap S_2$. Let $S_i = \{\triangle_{i,1}, \triangle_{i,2}, \dots, \triangle_{i,m}\}$ and $\triangle = \triangle_{1,j} = \triangle_{2,t}$ for some t, j. Then, by the above similar argument, $\{\triangle_{1,j-1}, \triangle_{1,j+1}\} = \{\triangle_{2,t-1}, \triangle_{2,t+1}\}$. Again, we consider $\triangle_{1,j-1}$ and repeat as above and continue. Hence, we get $S_1 = S_2$ after finite number of steps and the geometric carrier $|S_1|$ is bounded by identical cycle C. In this case, it defines a $K(r, 1, k)$ representation (this polyhedral representation is defined later in this section) of

M. (For an example in Figure 1, $K(7,4,1)$ is a representation of M where $s = 4$.) But, by Lemma 2.6, $K(r,1,k)$ does not exist. Therefore, $S_1 \cap S_2 = \emptyset$ and $S = S_1 \cup S_2$ where one of $|S_1|, |S_2|$ is a cylinder and another is a Möbius strip. Observe that both $|S_1|, |S_2|$ can not be Möbius strips as $S_1 \cap S_2 = \emptyset$. We denote Möbius strip by $\mathcal{M}_{\triangle}(C,M)$ which is bounded by C and $|S|$ by $S\mathcal{M}_{\triangle}(C,M)$. (For examples, $\mathcal{M}_{\triangle}(C,M)$ and $S\mathcal{M}_{\triangle}(C,M)$ are in Figure 5.) $\square$

Remark 2.1. Note that we use these above notions of $S(C,M), \mathcal{M}_{F_1,\ldots,F_t}(C,M)$ and $S\mathcal{M}_{F_1,\ldots,F_t}(C,M)$ in the subsequent sections. Here, the $S(C,M)$ used to denote a cylinder which is geometric career of the faces that are incident with C. The $\mathcal{M}_{F_1,\ldots,F_t}(C,M)$ used to denote a Möbius strip which is bounded by C and consists of faces of types $F_1, \ldots, F_t$. The $S\mathcal{M}_{F_1,\ldots,F_t}(C,M) := |S \cup \mathcal{M}|$ used to denote union of a cylinder $|S|$ and a Möbius strip $|\mathcal{M}|$ such that $|S| \cap |\mathcal{M}| = C$ and consists of faces of types $F_1, \ldots, F_t$. In this section, $t = 1$ and $F_1 = \triangle$.


v_2 v_1 v_7 v_6 v_5 v_4 v_3 v_2
u_1 u_2 u_3 u_4 u_5 u_6 u_7 u_1
x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_1
w_1 w_2 w_3 w_4 w_5 w_6 w_7 w_1
v_1 v_2 v_3 v_4 v_5 v_6 v_7 v_1

Figure 1 : M


u_5 u_6 u_7 u_1
u_1 u_2 u_3 u_4 u_5 u_6 u_7 u_1
x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_1
w_1 w_2 w_3 w_4 w_5 w_6 w_7 w_1
w_5 w_6 w_7 w_1

Figure 6 : $K(S'(M), \mathcal{M}', \mathcal{M}'')$


w_{r+1} w_{r+2} w_{r+3} w_{r+4} w_{2r-1} w_{2r} w_{2r+1} w_1 w_{r-2} w_{r-1} w_r w_{r+1}
w_{r+2} w_1
w_1 w_2 w_3 w_4 w_{r-1} w_r w_{r+1} w_{2r-1} w_{2r} w_{2r+1}
v_1 v_2 v_3 v_4 v_{r-1} v_r v_{r+1} v_{r+2} v_{2r-1} v_{2r} v_{2r+1} v_1

Figure 4


x_1 x_2 x_3 x_4 x_{r-2} x_{r-1} x_r x_1
w_1 w_2 w_3 w_4 w_{r-2} w_{r-1} w_r w_1
v_1 v_2 v_3 v_4 v_{r-2} v_{r-1} v_r v_1

Figure 2 : $S(C,M)$: Cylinder


w_{r+1} w_{r+2} w_{r+3} w_{r+4} w_{2r-1} w_{2r} w_{2r+1} w_1
w_{r+2} w_{2r-1} w_{2r} w_{2r+1} w_1
w_1 w_2 w_3 w_4 w_{r-1} w_r w_{r+1}
v_1 v_2 v_3 v_4 v_{r-1} v_r v_{r+1} v_{r+2} v_{2r-1} v_{2r} v_{2r+1} v_1

Figure 5 : $S\mathcal{M}_{\triangle}(C,M)$


x_1 x_2 x_3 x_4 x_{r-2} x_{r-1} x_r v_1
w_1 w_2 w_3 w_4 w_{r-2} w_{r-1} w_r w_1
v_1 v_2 v_3 v_4 v_{r-2} v_{r-1} v_r x_1

Figure 3

Let C be a cycle of type A in M. The $S(C,M)$ is a cylinder and let $\partial S(C,M) = \{C_1, C_2\}$. Observe as above that cycle C_1, C_2 are of type A and length(C) = length(C) = length(C_2). Let $C_1, C_2, \ldots, C_m$ be a sequence of cycles which are homologous to C in M. For some i, j, there is a cylinder in M which is bounded by C_i and C_j. So, C_i and C_j are homologous and have same length (see its similar argument in *Section* 2 [7]). That is, we get a cylinder $S_{l,t}$ for some cycles C_l and C_t such that $\partial S_{l,t} = \{C_t, C_l\}$. So, length$(C_t)$ = length(C_l). Thus, length(C_i) = length(C_j) $\forall\ i, j \in \{1, 2, \ldots, m\}$.

By Lemma 2.2, we assume that $S(C,M)$ contains in M. Here, $\partial S(C,M) = \{C', C''\}$ where C', C'' are two cycles of type A. Let $IF(L)$ denote a set of faces which are incident with L and $F(S(L,M))$ denote a set of faces of $S(L,M)$. At 1^{st} step, let $S = F(S(C,M)) \cup IF(C') \cup IF(C'')$ and $S_1 = S$. Next step, we consider S_1 in place of S and continue with as above. In this process, let S_i denote a set at i^{th} step and $\partial S_i = \{L_{i,1}, L_{i,2}\}$. Here,

$L_{i,1} \neq L_{i,2}$, $L_{i,1} = L_{i,2}$, $L_{i,1} = \partial\mathcal{M}'$, $L_{i,1} = \partial\mathcal{M}''$ or '$L_{i,1} = \partial\mathcal{M}'$ and $L_{i,2} = \partial\mathcal{M}''$' for some Möbius strips $\mathcal{M}', \mathcal{M}''$. This process stops if $L_{j,1} = L_{j,2}$ or '$L_{j,1} = \partial\mathcal{M}'$ and $L_{j,2} = \partial\mathcal{M}''$'. Let the above process stop at k^{th} step. Then, the $|S_k|$ is bounded by identical cycles or two Möbius strips. Hence, we get a cylindrical representation $S(M)$ of M which is bounded by two identical cycles, i.e., $L_{k,1} = L_{k,2}$ or a $S\mathcal{M}(M)$ which is bounded by two Möbius strips, i.e., $L_{k,1} = \partial\mathcal{M}'$ and $L_{k,2} = \partial\mathcal{M}''$. Again, by Lemma 2.2, we assume that $S\mathcal{M}_{\triangle}(C, M)$ contains in M. Here $\partial\mathcal{M}_{\triangle}(C, M)$ is a cycle. We repeat as above and consider $S\mathcal{M}_{\triangle}(C, M)$ in place of $S(C, M)$ in the above process. In this process, we get one boundary at each step. Thus, we get a $S\mathcal{M}(M)$ which is bounded by two Möbius strips. Therefore, by combining above two cases, we have

Lemma 2.3. *The map M has a $S(M)$ or a $S\mathcal{M}(M)$ representation.*

We first assume that $S(M)$ denotes M. So, $\partial S(M) = \{C, C'\}$ and $C = C'$.

Claim : The $S(M)$ has a (r, s, k)-representation ($K(r, s, k)$ representation) for some $r, s, k \in \mathbb{N}$.

We recall similar definition of (r, s, k)-representation as in *Section* 2 [7]. Let $v \in V(C)$. By definition of A, we have three paths of type A through v. Let L_1, L_2, L_3 denote three paths of type A through v in $S(M)$. Let $L_1 = P(a_1, a_2, \ldots, a_r)$ and $C = L_1$. We make a cut starting from v along $P = P(a_1 = w_1, \ldots, w_s = a_{k+1}) \subset L_3$ until reaching C' again for the first time. Let s denote the number of cycles which are homologous to C along P. Observer that the number s is equal to the length of the path P, that is, $s = m$. So, length$(C) = r$ and s is the number of horizontal cycles. Thus, we get a planar polyhedral representation of M and denote it by (r, s)*-representation*. We say that cycle C is the *lower (base) horizontal cycle* [7] and the other cycle C' is *upper horizontal cycle* [7] in (r, s)-representation. The (r, s)-representation has identification of vertical sides (non homologous to the horizontal cycles) in the natural manner but the identification of the horizontal sides needs some shifting with twist so that a vertex in the lower(base) side is identified with a vertex in the upper side. Let $C' = C(a_{k+1}(= w_m), a_k, a_{k-1}, \ldots, a_k)$. Then, the vertex a_{k+1} is the starting vertex of the upper horizontal cycle C' in (r, s)-representation. The $k = \text{length}(P(a_1, \ldots, a_{k+1}))$ where $P(a_1, \ldots, a_{k+1}) \subset C$. So, we denote the (r, s)-representation by (r, s, k)*-representation* where r, s, k are defined as above. We say that boundaries of (r, s, k)-representation are the cycles and paths along which we took the cuts to construct (r, s, k)-representation. By this construction, a (r, s, k)-representation exists in $S(M)$. In this article, we denote (r, s, k)-representation by $K(r, s, k)$. (For example in Figure 1, $K(7, 4, 1)$ denote a $(7, 4, 1)$-representation, $C = C(v_1, \ldots, v_7), C' = C(v_2, v_1, v_7, v_6, v_5, v_4, v_3), P = P(v_1, w_1, x_1, u_1, v_2)$.) It's clear from the definition that $S(M)$ has a (r, s, k)-representation. This proves the claim.

Next we assume that $S\mathcal{M}(M)$ denotes M.

Claim : The $S\mathcal{M}(M)$ has $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation.

The $S\mathcal{M}(M)$ bounded by two Möbius strips. We cut along the boundaries of the Möbius strips in $S\mathcal{M}(M)$. In this case, we get three components. These are a cylinder namely $S'(M)$ and two Möbius strips namely $\mathcal{M}', \mathcal{M}''$ where $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}_{\triangle}(L, M)\}$ for some cycle L of type A. We denote this representation by $K(S'(M), \mathcal{M}', \mathcal{M}'')$. This proves the claim. (For an example in Figure 6, $K(S'(M), \mathcal{M}', \mathcal{M}'')$ denotes a representation of $M, S'(M) = |\{x_1x_2w_1, w_1w_2x_2, x_2x_3w_2, w_2w_3x_3, x_3x_4w_3, w_3w_4x_4, x_4x_5w_4, w_4w_5x_5, x_5x_6w_5, w_5w_6x_6, x_6x_7w_6, w_6w_7x_7, x_7x_1w_7, w_7w_1x_1, u_1u_2x_1, x_1x_2u_2, u_2u_3x_2, x_2x_3u_3, u_3u_4x_3, x_3x_4u_4, u_4u_5x_4, x_4x_5u_5, u_5u_6x_5, x_5x_6u_6, u_6u_7x_6, x_6x_7u_7, u_7u_1x_7, x_7x_1u_1\}|, \mathcal{M}' = |\{u_4u_5u_1, u_1u_2u_5,$

$u_5u_6u_2,\ u_2u_3u_6, u_6u_7u_3, u_3u_4u_7, u_7u_1u_4\}|$ and $\mathcal{M}'' = |\{w_1w_2w_5, w_5w_6w_2, w_2w_3w_6, w_6w_7w_3, w_3w_4w_7, w_7w_1w_4, w_4w_5w_1\}|$.)

So, the map M has a $K(r,s,k)$ representation or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation. We first assume that $K(r,s,k)$ represents M. Let L, L' denote the lower and upper horizontal cycle respectively in $K(r,s,k)$. Let b be a vertex in L. By definition of A, let L_1, L_2, L_3 denote three paths of type A through b. Let $L_1 = L = L', L_2 = P(b, \ldots, x)$ and $L_3 = P(b, \ldots, y)$ where $x, y \in V(L)$. The paths L_2, L_3 are vertical and non-homologous to L in $K(r,s,k)$. Let triangles $\triangle'_1, \triangle'_2, \ldots, \triangle'_m$ lie on one side of L_2 and $\triangle''_1, \triangle''_2, \ldots, \triangle''_m$ lie on the other side of L_2 in $K(r,s,k)$. Observe that the geometric career $|\{\triangle'_1, \triangle'_2, \ldots, \triangle'_m\}|$ is bounded by $L_2, e', L_{2,1} := P(b', \ldots, x')$, and e where $e = bb' \in E(L)$ and $e' = xx' \in E(L')$ in $K(r,s,k)$. Similarly, the geometric career $|\{\triangle''_1, \triangle''_2, \ldots, \triangle''_m\}|$ is bounded by $L_2, \bar{e}', L_{2,2}$, and $\bar{e}$ where $\bar{e} \in E(L)$ and $\bar{e}' \in E(L')$ in $K(r,s,k)$. Hence, we get two paths $L_{2,1}$ and $L_{2,2}$. Again, we consider $L_{2,2}$ in place of L_2 and repeat as above and continue. Hence, we get a sequence of paths $L_{2,1}, L_{2,2}, \ldots, L_{2,s}$. Similarly, we repeat with above process and consider cycle L_3 in place of L_2. Hence, we get an another sequence of paths namely $L_{3,1}, L_{3,2} \ldots, L_{3,s}$. Let $L_{2,1} = P(b', \ldots, x')$. We extend $L_{2,1}$ along the path of type A. Since $L = L'$ and have opposite orientations, so, by definition of A, after the vertex x' the path $L_{2,1}$ extends along a path $L_{3,j}$ for some j. Here, we repeat this process for each $L_{2,i}$ where $1 \le i \le s$. Let $L_{2,i}$ extends along $L_{3,j}$ for some j. So, we get a path $L_{2,i} \cup L_{3,j}$. Thus, we get a sequence of paths $R_1 := L_{2,1} \cup L_{3,i_1}, R_2 := L_{2,2} \cup L_{3,i_2}, \ldots, R_s := L_{2,s} \cup L_{3,i_s}$. So, we have three paths $L, L_{2,i}$ and $L_{3,j}$ through b for some i, j. The paths $L_{2,i}$ and $L_{3,j}$ are part of some R_{l_1} and R_{l_2}. Therefore, let P', P'', P''' be three paths of type A through $b \in V(L)$. Then, $P' \subset R_k, P'' \subset R_{k'}$ and $P''' \subset L$ for some k, k'. Since the paths R_k and $R_{k'}$ are same type and part of some walk of type A in M, it follows that we have only one cycle L of type A through b. Therefore, we have only one cycle of type A through each vertex of M. Combining above all cases, we have

Lemma 2.4. *The map M has a $K(r,s,k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation but not both.*

Proof. By the preceding section, the map M has either a $K(r,s,k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation. In a map M, we have one cycle of type A as above though each vertex of M. We are using this cycle to construct both the representations. Since there is only one cycle of type A through a vertex and we are using this cycle to construct the above two representations, it implies that map M can not have both the representations simultaneously. So, $K(r,s,k)$ and $K(S'(M), \mathcal{M}', \mathcal{M}'')$ are non isomorphic. That is, M does not have both the representations. Therefore, we study both the representations bellow separately. This completes the proof. □

Lemma 2.5. *The cycles of type A in M have same length.*

Proof. Let $v \in V(M)$. The M has either a $K(r,s,k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation. In $K(r,s,k)$, we have one cycle and one walk through v as above. Let $C'_1, \ldots, C'_s$ denote a sequence of horizontal cycles of type A which are homologous to C'_1 in $K(r,s,k)$. By the preceding argument, length(C'_i) = length(C'_j) for all $1 \le i, j \le s$. So, there is an unique cycle of type A in $K(r,s,k)$ upto homologous. Therefore, the cycles of type A have same length. Similarly we can proceed with argument as above in $K(S'(M), \mathcal{M}', \mathcal{M}'')$ and hence, we get same result. This completes the proof. □

In Lemma 2.4, the representations $K(r,s,k)$ and $K(S'(M),\mathcal{M}',\mathcal{M}'')$ are non-isomorphic and these representations are exist in maps of type $\{3^6\}$ on the torus. So, we study both the representations separately and classify them in bellow.

Classification of $K(r,s,k)$ on n vertices : We define admissible relations among r,s,k of $K(r,s,k)$ such that $K(r,s,k)$ represents a map after identifying its boundaries.

Lemma 2.6. *The maps of type $\{3^6\}$ of the form $K(r,s,k)$ exist if and only if the following holds : (i) $rs \geq 9$, (ii) $r \geq 3$, (iii) $s \geq 3$ and (iv) $k \in \{t : 0 \leq t \leq r-1\}$ if $s \geq 3$.*

Proof. One can repeat similar arguments as in *Section* 2 [7] and study each cases. If $s \leq 2$ or $r \leq 2$ or $\text{rs} < 9$ in $K(r,s,k)$ then we get some vertex whose link is not a cycle. Similarly, it is easy to observe that $k \in \{t : 0 \leq t \leq r-1\}$ if $s \geq 3$ in $K(r,s,k)$ since length of the lower horizontal cycle is r. □

Let M_1 and M_2 be two maps of type $\{3^6\}$ on the Klein bottle. Let $K(r_i,s_i,k_i)$ denote a planar polyhedral representation of M_i. Now we have

Lemma 2.7. *Let $K(r_i,s_i,k_i)$ denote a (r_i,s_i,k_i) -representation of M_i on n vertices for $i \in \{1,2\}$. Then, map $M_1 \cong M_2$ if $(r_1,s_1,l_1) = (r_2,s_2,l_2)$ where $l_i \in \{k_i \; mod(2), (k_i + r_i) \; mod(2)\}$.*

Proof. First assume that $r_1 = r_2, s_1 = s_2, k_1 = k_2$. Let $C(1,0), C(1,1), \ldots C(1,s_1-1)$ denote a sequence of horizontal cycles of type A in $K(r_1,s_1,k_1)$. Define $C(1,0) := C(u_{0,0},u_{0,1},\ldots,u_{0,r_1-1}), C(1,1) := C(u_{1,0},u_{1,1},\ldots,u_{1,r_1-1}),\ldots,\ C(1,s_1-1) := C(u_{s_1-1,0}, u_{s_1-1,1},\ldots,u_{s_1-1,r_1-1})$. Again, let $C(2,0), C(2,1), \ldots C(2,s_2-1)$ denote a sequence of horizontal cycles of type A in $K(r_2,s_2,k_2)$. Define $C(2,0) := C(v_{0,0},v_{0,1},\ldots,v_{0,r_2-1}), C(2,1) := C(v_{1,0},v_{1,1},\ldots,v_{1,r_2-1},v_{1,0}),\ldots,\ C(2,s_2-1) := C(v_{s_2-1,0},v_{s_2-1,1},\ldots,v_{s_2-1,r_2-1})$. Since maps of type $\{3^6\}$ contain unique cycle of type A up to homologous, it follows by Lemma 2.6 that $r_1 = r_2$ and $s_1 = s_2$. So, we have the following cases.

Case 1 : If $k = k_1 = k_2$ then define a map $f_1 : V(K(r_1,s_1,k_1)) \to V(K(r_2,s_2,\ k_2))$ by $f_1(u_{t,i}) = v_{t,i}$ for $0 \leq t \leq s-1$ and $0 \leq i \leq r-1$. By definition, $f_1(lk(u_{t,i})) = f_1(C(u_{t-1,i-1},\ u_{t-1,i},\ u_{t-1,i+1},\ u_{t,i+1},\ u_{t+1,i+1},\ u_{t+1,i},\ u_{t,i-1})) = (f_1(u_{t-1,i-1}),\ f_1(u_{t-1,i}),\ f_1(u_{t-1,i+1}),\ f_1(u_{t,i+1}),\ f_1(u_{t+1,i+1}),\ f_1(u_{t+1,i}),\ f_1(u_{t,i-1})) = C(v_{t-1,i-1},\ v_{t-1,i},\ v_{t-1,i+1},\ v_{t,i+1},\ v_{t+1,i+1},\ v_{t+1,i}, v_{t,i-1})$. So, $f_1(lk(u_{t,i})) = lk(v_{t,i})$. That is, f_1 sends vertices to vertices, edges to edges, faces to faces and also, preserves incidents. So, f_1 is an isomorphism map. Therefore, $K(r_1,s_1,k_1) \cong K(r_2,s_2,k_2)$.

Case 2 : If $k_1 \neq k_2$ then by assumption, let $l_1 = l_2 = 0$. Here, we have the following cases : $2 \mid k_i$ or $2 \mid (k_i + r_i)$.

Let $2 \mid k_i$. The vertex $u_{0,\frac{k_i}{2}} \in V(C(i,0))$. Let Q_1, Q_2, Q_3 denote three paths of type A through $u_{0,\frac{k_i}{2}}$ in $K(r_i,s_i,k_i)$. Let $C(i,s_i) := C(i,0), Q_1 = C(i,0)$ and $Q_3 = P(u_{0,\frac{k_i}{2}},\ \ldots,\ w(= u_{0,\frac{k_i}{2}}))$ where $w \in V(C(i,s_i))$. We identify $K(r_i,s_i,k_i)$ along vertical boundaries and cut along Q_3. In this process, we are not changing the horizontal cycles and number of horizontal cycles vertically. So in this case, we get a $K(r_i,s_i,0)$ representation of M_i.

Let $2 \mid (k_i + r_i)$. Similarly as above, $u_{0,\frac{k_i+r_i}{2}} \in V(C(i,0))$. Let L_1, L_2, L_3 denote three paths through $u_{0,\frac{k_i+r_i}{2}}$ in $T(r_i,s_i,k_i)$. Let $C(i,s_i) := C(i,0), L_1 = C(i,0)$ and $L_3 = P(u_{0,\frac{k_i+r_i}{2}},\ldots,v(= u_{0,\frac{k_i+r_i}{2}}))$ where $v \in V(C(i,s_i))$. We identify $T(r_i,s_i,k_i)$ along vertical boundaries and cut along L_3. Hence, we get a $K(r_i,s_i,0)$ representation of M_i.

So, we get a $K(r_i,s_i,0)$ representation from $K(r_i,s_i,k_i)$ if $2 \mid k_i$ or $2 \mid (k_i + r_i)$. Since $r_1 = r_2$ and $s_1 = s_2$, it follows by $f_1, K(r_1,s_1,0) \cong K(r_2,s_2,0)$.

Again, let $l_1 = l_2 = 1$. If $2 \mid (k_i - 1)$ then similarly as above, we identify $K(r_i, s_i, k_i)$ along their vertical boundaries and cut along the vertical path of type A through $u_{0,\frac{k_i-1}{2}} \in V(C(i,0))$ in $K(r_i, s_i, k_i)$. Again, if $2 \nmid (k_i - 1)$ then $2 \mid (k_i + r_i - 1)$. In this case, we consider vertical path of type A through $u_{0,\frac{k_i+r_i-1}{2}} \in V(C(i,0))$. Here, in both the cases, we get a $K(r_i, s_i, 1)$ representation of M_i. Since $r_1 = r_2$ and $s_1 = s_2$, if follows by $f_1, K(r_1, s_1, 1) \cong K(r_2, s_2, 1)$.

Hence by f_1, $M_1 \cong M_2$. This completes the proof. ☐

Thus, we have

Corollary 2.1. *Let $K(r_i, s_i, k_i)$ denote a (r_i, s_i, k_i)-representation of M_i on n vertices for $i \in \{1, 2\}$. Then, $K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2)$ $\forall$ $r_1 \neq r_2, K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2)$ $\forall$ $s_1 \neq s_2, K(r_i, s_i, 0) \cong K(r_i, s_i, k_i)$ for $2 \mid k_i$ or $2 \mid (k_i + r_i)$, and $K(r_i, s_i, 1) \cong K(r_i, s_i, k_i)$ for $2 \mid (k_i - 1)$ or $2 \mid (k_i + r_i - 1)$.*

Classification of $K(l, t)$ on n vertices : Let $S\mathcal{M}(M)$ denote a representation of M on n vertices. By Lemma 2.4, the $S\mathcal{M}(M)$ has a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation where $C' = \partial\mathcal{M}', C'' = \partial\mathcal{M}''$ are two homolgous cycles of type A and $\partial S'(M) = \{C', C''\}$. Hence, we denote $K(S'(M), \mathcal{M}', \mathcal{M}'')$ by $K(l, t)$ if $l = length(C') = length(C'')$ and t denote the number of cycles of type A in $S'(M)$.

Lemma 2.8. *The maps of type $\{3^6\}$ of the form $K(l, t)$ exist if and only if the following holds : (i) $tl \geq 10$, (ii) $t \geq 2$, (iii) $2 \nmid l$ and $l \geq 5$.*

Proof. If $t = 1$ then we get some vertex in C' whose link is not a cycle. So, $S'(M)$ contains at least two disjoint cycles of type A and hence, $t \geq 2$. Again, the $\mathcal{M}'$ is a Möbius strip which has one boundary. By the argument as in Lemma 2.2, $\partial\mathcal{M}' \subset \partial S'(M)$. Let t_1 denote the number of cycles of type A in $\mathcal{M}'$. Then, by the argument of Lemma 2.2, $t_1 = 1$. Similarly, let t_2 denote the number of cycles of type A in $\mathcal{M}''$ then $t_2 = 1$. Therefore, $(t + t_1 + t_2 - 2) \geq 2$ as $\partial S'(M) = \{\partial\mathcal{M}', \partial\mathcal{M}''\}$. The cycles of type A covers all the vertices of the map M. So, $n = (t + t_1 + t_2 - 2)l = tl$. This implies that $tl \mid n, l \mid n$ and $t \mid n$. By the argument of Lemma 2.2, $2 \nmid l$. We follow similar argument as in Lemma 2.6 and hence, $l \geq 5$ and $tl \geq 10$. This completes the proof. ☐

Let $K(l_i, t_i)$ denote a representation of M_i on n vertices. Then, we have

Lemma 2.9. *The $K(l_1, t_1) \cong K(l_2, t_2)$ if $l_1 = l_2$.*

Proof. Let $l_1 = l_2$. Let $C_1, \ldots, C_s$ denote cycles of type A in M_1. Let $C_1 = \partial\mathcal{M}'_1$ and $C_s = \partial\mathcal{M}''_1$. Again, let $L_1, \ldots, L_s$ denote cycles of type A in M_2. Let $L_1 = \partial\mathcal{M}'_2$ and $L_s = \partial\mathcal{M}''_2$. We follow similar argument as in Lemma 2.7, and hence, we define an isomorphism map $f_1 : V(M_1) \to V(M_2)$ by $f_1(V(C_i)) = V(L_i)$ for $1 \leq i \leq s$. So, by $f_1, K(l_1, t_1) \cong K(l_2, t_2)$ i.e. $M_1 \cong M_2$. This completes the proof. ☐

Thus, we have

Corollary 2.2. *Let $K(l_i, t_i)$ denote $K(l_i, S'(M_i), \mathcal{M}'_i, \mathcal{M}''_i)$ representation of M_i on n vertices for $i \in \{1, 2\}$. Then, $K(l_1, t_1) \ncong K(l_2, t_2)$ $\forall$ $l_1 \neq l_2$ and $K(l_1, t_1) \ncong K(l_2, t_2)$ $\forall$ $t_1 \neq t_2$.*

2.2 Maps of type $\{4^4\}$: Let M be a map of type $\{4^4\}$ on the Klein bottle. A path $P(\dots, u_{i-1},\, u_i,\, u_{i+1}, \dots)$ in edge graph of M is of type B if $lk(u_i) = C(\mathbf{a}, b, \mathbf{c}u_{i+1}, \mathbf{d}, e, \mathbf{f}, u_{i-1})$ implies $lk(u_{i+1}) = C(\mathbf{b}, c, \mathbf{g}u_{i+2}, \mathbf{h}, d, \mathbf{e}, u_i)$ and $lk(u_{i-1}) = C(\mathbf{k}, a, \mathbf{b}u_i, \mathbf{e}, f, \mathbf{l}, u_{i-2})$ at u_i for all i. In the definition, the bold vertex represents non adjacent vertex. Let W be a maximal walk of type B. Then, similarly as in Section 2.1, it is either a cycle or a walk. Since degree of a vertex is four, it implies that every closed walk of type B is a cycle. Let C be a cycle of type B in M and S denote a set of faces which are incident with C. We argue similarly as in Lemma 2.2. In the proof we consider quadrangles in place of triangles and repeat similar argument as in Lemma 2.2. Hence, we get a geometric carrier $|S|$ which is either $S(C,M), \mathcal{M}(C,M)$ or $S\mathcal{M}_4(C,M)$. So, the map M contains $S(C,M), \mathcal{M}(C,M)$ or $S\mathcal{M}_4(C,M)$ for some cycle C of type B in M. (see Figure 7, it is an example of a Möbius strip $\mathcal{M}(C,M)$ where $C = C(w_1, \dots, w_r)$.) We consider each of $S(C,M), \mathcal{M}(C,M)$ and $\mathcal{M}_4(C,M)$ and repeat the similar arguments as in Lemma 2.3. Thus, we get a representation of M which is either a $S(M)$ which is bounded by identical cycles or a $S\mathcal{M}(M)$ with two Möbius strips $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}(C,M), \mathcal{M}_4(L,M) \mid$ for some cycles L, C of type $B\}$. Therefore, the M has $S(M)$ or $S\mathcal{M}(M)$ representation.


u_4 u_5 u_6 u_1
u_1 u_2 u_3 u_4 u_5 u_6 u_1
x_1 x_2 x_3 x_4 x_5 x_6 x_1
w_1 w_2 w_3 w_4 w_5 w_6 w_1
w_4 w_5 w_6 w_1

Figure 8


x_1 x_2 x_3 x_4 x_{r-2} x_{r-1} x_r v_1
w_1 w_2 w_3 w_4 w_{r-2} w_{r-1} w_r w_1
v_1 v_2 v_3 v_4 v_{r-2} v_{r-1} v_r x_1

Figure 7 : $\mathcal{M}(C, M)$


w_1 w_4 x_4 u_4 u_1 x_1 w_1
w_6 w_3 x_3 u_3 u_6 x_6 w_6
w_5 w_2 x_2 u_2 u_5 x_5 w_5
w_4 w_1 x_1 u_1 u_4 x_4 w_4

Figure 9 : K(6, 3, 1)

Let $S(M)$ denote a representation of M. Let $\partial S(M) = \{C, C'\}$ where C, C' are two cycles of type B. We repeat the similar argument as in Section 2.1 which is done to define (r,s,k)-representation of a map. Let $v \in V(C)$ and $L = P(v, \dots, w)$ be a path of type B where $w \in V(C')$ and L is not homologous to C. Then, we take second cut along L. Hence, we get a (r,s,k)-representation of M and it is denoted by $K(r,s,k)$ for some r,s,k. Let $S\mathcal{M}(M)$ denote a representation of M. We cut along the boundaries of Möbius strips, and hence, we get three components which are a cylinder namely $S'(M)$ and two Möbius strips namely $\mathcal{M}', \mathcal{M}''$ where $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}(L_1, M), \mathcal{M}_4(L_2, M) \mid L_1, L_2$ are two cycles of type $B\}$. Therefore, by the similar argument as in Section 2.1, the map M has a $K(r,s,k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation. So, we have

Lemma 2.10. *The map M has a $K(r,s,k)$ representation.*

Proof. By the preceding above arguments, the map M has either a $K(r,s,k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation. Assume that $K(S'(M), \mathcal{M}', \mathcal{M}'')$ denotes M. Let $\partial S'(M) = \{C, C'\}, C = C(u_1, \dots, u_l)$ and $C' = C(v_1, \dots, v_l)$, and $\mathcal{M}' = \mathcal{M}(L, M), \mathcal{M}'' = \mathcal{M}(L', M)$ for some cycles L, L' of type B. (For an example in Figure 8, $L = C(u_1, \dots, u_6), L' = C(w_1, w_2, \dots, w_6), \mathcal{M}' = \mathcal{M}_4(L, M) = |\{[u_1, u_2, u_5, u_4], [u_2, u_3, u_6, u_5], [u_3, u_4, u_1, u_6]\}|, \mathcal{M}'' = \mathcal{M}_4(L', M) = |\{[w_1, w_2, w_5, w_4], [w_2, w_3, w_6, w_5], [w_3, w_4, w_1, w_6]\}|$ and $S'(M) = |\{[x_1, x_2, w_2, w_1], [x_2, x_3, w_3, w_2], [x_3, x_4, w_4, w_3], [x_4, x_5, w_5, w_4], [x_5, x_6, w_6, w_5], [x_6, x_1, w_1, w_6], [x_1, x_2, u_2, u_1], [x_2, x_3, u_3, u_2], [x_3, x_4, u_4, u_3], [x_4, x_5, u_5, u_4], [x_5, x_6, u_6, u_5], [x_6, x_1, u_1, u_6]\}|$.) Let $u \in V(L)$. We cut $\mathcal{M}(L, M)$ along the path of type B through u. Thus, we get a rectangular planar representation of $\mathcal{M}(L, M)$ which has three horizontal paths of same length. Since above and lower paths define a cycle, it implies that its length is twice of length(L). By definition of B, we have two paths Q_1, Q_2 of type B through u_1. Let $Q_1 \subset C$. In this case, we cut $\mathcal{M}(L, M)$ along the path $P(u_1, x, u_{l'+1}) \subset Q_2$ of type B where $x \in V(L)$. Hence, we get a planar polyhedral representation, say $\square(\mathcal{M})$. (For an example in Figure 8, we cut $\mathcal{M}(L, M)$ along $P(u_1, u_4)$ and hence, we get $\square(\mathcal{M}_4)$ which is bounded by $P(u_1, u_4), P(u_4, u_5, u_6, u_1), P(u_1, u_4), P(u_4, u_3, u_2, u_1)$.) Again, let Z_1, Z_2 denote two paths of type B through $u_{l'+1}$. Let $Z_1 \subset C$. Similarly, we cut $S'(M)$ along the path $P(u_1, \ldots, v_t) \subset Q_2$ of type B and then, along $P(u_{l'+1}, \ldots, v_{l'+t}) \subset Z_2$ of type B. Hence, we get two planar polyhedral representations $\square'$ and $\square''$. (For example in Figure 8, we cut $S'(M)$ along $P(u_1, x_1, w_1)$ and $P(u_4, x_4, w_4)$ and we get $\square' = |\{[x_1, x_2, w_2, w_1], [x_2, x_3, w_3, w_2], [x_3, x_4, w_4, w_3], [x_1, x_2, u_2, u_1], [x_2, x_3, u_3, u_2], [x_3, x_4, u_4, u_3]\}|$ and $\square'' = |\{[x_4, x_5, u_5, u_4], [x_5, x_6, u_6, u_5], [x_6, x_1, u_1, u_6], [x_4, x_5, w_5, w_4], [x_5, x_6, w_6, w_5], [x_6, x_1, w_1, w_6]\}|$.) Similarly, we cut $\mathcal{M}(L', M)$ along $P(v_t, z, v_{l'+t}) \subset Q_2$ of type B where $z \in V(L')$. Hence, we get $\square(\mathcal{M}')$. (For example in Figure 8, we cut $\mathcal{M}(L', M)$ along $P(w_1, w_4)$ and hence, we get $\square(\mathcal{M}_4')$ which is bounded by $P(w_1, w_4), P(w_4, w_5, w_6, w_1), P(w_1, w_4), P(w_4, w_3, w_2, w_1)$.) The boundaries of $\square(\mathcal{M})$ are $P(u_1, x, u_{l'+1}), P(u_{l'+1}, \ldots, u_l, u_1), P(u_{l'+1}, x, u_1)$ and $P(u_{l'+1}, \ldots, u_1)$. The boundaries of $\square'$ are $P(u_1, u_2, \ldots, u_{l'+1}), P(u_{l'+1}, \ldots, v_{t+l'}), P(v_{t+l'}, \ldots, v_t)$ and $P(v_t, \ldots, u_2, u_1)$. The boundaries of $\square''$ are $P(u_{l'+1}, \ldots, u_l, u_1), P(u_1, \ldots, v_t), P(v_t, \ldots, v_{t+l'})$ and $P(v_{t+l'}, \ldots, u_{l'+1})$. The boundaries of $\square(\mathcal{M}')$ are $P(v_t, z, v_{l'+t}), P(v_{l'+t}, \ldots, v_t), P(v_t, z, v_{t+l'})$ and $P(v_{l'+t}, \ldots, v_t)$. Now, we consider $\square(\mathcal{M})$ and $\square'$, and identify along $P(u_1, \ldots, u_{l'+1})$. Then, we consider $\square(\mathcal{M})$ and $\square''$, and identify along $P(u_{l'+1}, \ldots, u_1)$. Again, we consider $\square''$ and $\square(\mathcal{M}')$, and identify along $P(v_{t+l'}, \ldots, v_t)$. Hence, we get a planar polyhedral representation of $K(S'(M), \mathcal{M}', \mathcal{M}'')$ which is bounded by cycle $\mathcal{C} = P(v_t, \ldots, u_1) \cup P(u_1, \ldots, u_{l'+1}) \cup P(u_{l'+1}, \ldots, v_{t+l'}) \cup P(v_{t+l'}, \ldots, v_t)$ and path $P(v_t, \ldots, v_{t+l'})$. Hence, we get a $K(r', s', k')$ representation from $K(S'(M), \mathcal{M}', \mathcal{M}'')$ for some k' where $r' = \text{length}(\mathcal{C})$. (For example in Figure 9, we get $K(6, 3, 1)$ from Figure 8.) So, every map M has $K(r, s, k)$ representation for some r, s, k. Similar argument as above we repeat if '$\mathcal{M}' = \mathcal{M}_4(L, M)$ and $\mathcal{M}'' = \mathcal{M}(L', M)$', and if '$\mathcal{M}' = \mathcal{M}_4(L, M)$ and $\mathcal{M}'' = \mathcal{M}_4(L', M)$'. $\square$

So, by the above Lemma 2.10, $K(r, s, k)$ denotes M for some r, s, k. The $K(r, s, k)$ has two cycles of type B through each vertex. So, by the similar argument as in Lemma 2.5, it has two cycles of type B upto homologous. Let C' and C'' denote two cycles in M through a vertex. The cycles are non-contractible and represent two generators of the fundamental group $\mathbb{Z} \times \mathbb{Z}_2$ of the Klein bottle as it is discussed in Section 2.1. Since $\mathbb{Z} \ncong \mathbb{Z}_2$, it follows that through each vertex of M, there is only one cycle of type B which is homologous to $\mathbb{Z}$ and denote it by C'. In this case, by the above preceding argument, we get a $K(r, s, k)$ representation of M. Similarly, when we consider C'' then we get a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation of M. Since these two cycles represent generator of two different groups $\mathbb{Z}, \mathbb{Z}_2$, it implies that there is no isomorphism map which maps cycle C' to C''. Therefore, we consider cycle C' and its homologous cycles in the proof of Lemma 2.12. By the above Lemma 2.10, every map has a $K(r, s, k)$ representation. So, we study only $K(r, s, k)$ representations.

We define admissible relations among r, s, k of $K(r, s, k)$ such that $K(r, s, k)$ represents a map after identifying its boundaries. The proof of next Lemma 2.11 similar as in Lemma 2.6. We consider each case separately and if we consider value of r, s, k not belong to the mentioned ranges in Lemma 2.11 then we get some vertex whose link is not a cycle. Thus, we have

Lemma 2.11. *The maps of type $\{4^4\}$ of the form $K(r, s, k)$ exist if and only if the following holds : (i) $rs \geq 9$, (ii) $r \geq 3$, (iii) $s \geq 3$, and (iv) $k \in \{t : 0 \leq t \leq r-1\}$ if $s \geq 3$.*

Let M_1 and M_2 be two maps of type $\{4^4\}$ on the Klein bottle. Then, we have

Lemma 2.12. *Let $K(r_i, s_i, k_i)$ denote a (r_i, s_i, k_i) -representation of M_i on n vertices for $i \in \{1,2\}$. Then, $M_1 \cong M_2$ if $(r_1, s_1, l_1) = (r_2, s_2, l_2)$ where $l_i \in \{k_i \ mod(2), (k_i + r_i) \ mod(2)\}$.*

Proof. The proof repeats the similar argument as in Lemma 2.7. Let $r_1 = r_2, s_1 = s_2, k_1 = k_2$. Define an isomorphism map $f_1 : V(K(r_1, s_1, k_1)) \to V(K(r_2, s_2, k_2))$ as in Lemma 2.7. So, $M_1 \cong M_2$ by f_1. When $k_1 \neq k_2$ then we repeat similar argument as in Lemma 2.7. Hence, we get either a $K(r_i, s_i, 0)$ or a $K(r_i, s_i, 1)$ representation from $K(r_i, s_i, k_i)$. So, by $f_1, M_1 \cong M_2$. There are two cycles $L_{1,i}$ and $L_{2,i}$ through a vertex in M_i. Let $L_{1,i}$ denote a horizontal cycle in $K(r_i, s_i, k_i)$. By the above preceding argument, there is no isomorphism map which maps $L_{1,i}$ to $L_{2,i}$ since these two cycles represent generator of non-isomorphic groups. So, we do not consider cycle $L_{2,i}$ in the above case. Therefore, the given conditions are the sufficient conditions for isomorphism. This completes the proof. ☐

Thus, we have

Corollary 2.3. *Let $K(r_i, s_i, k_i)$ denote a (r_i, s_i, k_i)-representation of M_i on n vertices for $i \in \{1,2\}$. Then, $K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2) \ \forall \ r_1 \neq r_2, K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2) \ \forall \ s_1 \neq s_2, K(r_1, s_1, 0) \cong K(r_1, s_1, k_1)$ for $2 \mid k_1$ or $2 \mid (k_1 + r_1)$, and $K(r_1, s_1, 1) \cong K(r_1, s_1, k_1)$ for $2 \mid (k_1 - 1)$ or $2 \mid (k_1 + r_1 - 1)$.*

2.3 Maps of type $\{6^3\}$: In Section 2.1, we have shown that the maps of type $\{3^6\}$ on the Klein bottle can be classified up to isomorphism. Let $P_1, \ldots, P_k$ denote maps of type $\{3^6\}$ on n vertex. Let M_i denote the dual of P_i. Let $m = \#F(P_i)$. By the notion of duality, then $M_1, \ldots, M_k$ are the only maps of type $\{6^3\}$ with m vertices upto isomorphism. In [7], the details of the above claim has been discussed elaborately. Therefore, one can classify maps of type $\{6^3\}$ on the Klein bottle using the classification of maps of type $\{3^6\}$ on the Klein bottle. Thus, we have

Lemma 2.13. *The maps of type $\{6^3\}$ on the Klein bottle can be classified up to isomorphism using the classifcation of maps of type $\{3^6\}$ on the Klein bottle.*

2.4 Maps of type $\{3^3, 4^2\}$: Let M be a map of type $\{3^3, 4^2\}$ on the Klein bottle. We recall some definitions and results from Maity and Upadhyay [7] which are defined on the torus. These are also well defined (clear from the definitions) in the map on the Klein bottle. The definitions are as follows. Let $P(\ldots, u_{i-1}, u_i, u_{i+1}, \ldots)$ be a path in edge graph of M. The path P is of type A_1 [7] if all the triangles incident with u_i lie on one side and all quadrangles incident with u_i lie on the other side of P at u_i for all i. Let W be a maximal walk of type A_1 in M. From definition, it follows that there is only one path of type A_1 through each vertex of M. So, by the similar argument as in Lemma 2.1, the maximal walk W of type A_1 is a cycle of type A_1. Let C be a cycle of type A_1 in M. Let S denote a set of faces which are incident with C. Then, we have

Lemma 2.14. *The geometric carrier $|S|$ is $S(C, M), S\mathcal{M}_{\triangle}(C, M)$ or $S\mathcal{M}_{\square}(C, M)$.*

Proof. We proceed with the similar argument as in Lemma 2.2. Let $lk(w_1) = C(x_1, w_r, \boldsymbol{v}_r, v_1, \boldsymbol{v}_2, w_2, x_2)$. We consider $|S|$ and cut along the path $P(x_1, w_1, v_1)$. Hence, we have the following cases. If a path $P(x_1, w_1, v_1)$ identifies with $P(x_1, w_1, v_1)$ with out any twist then we get a cylinder (see an example in Figure 10, $C(w_1, \ldots, w_r)$). We denote it by $S(C, M)$. If $P(x_1, w_1, v_1)$ identifies with $P(x_1, w_1, v_1)$ with a twist then we get a Möbius strip (see an example in Figure 12). In this case, we consider $lk(w_1)$ and which is of type $\{3^2,4,3,4\}$. So, this case is not possible. Now by definition of A_1, the triangles which are incident with C are lie on one side of C and quadrangles are lie on the other side of C. Assume that the cycle $C=C(w_1,w_2,\ldots,w_{2r+1})$ is of odd length. By the similar argument as in Lemma 2.2, the triangles which are incident with C forms a Möbius strip $\mathcal{M}_{\triangle}(C,M)$. In this case, we get $S\mathcal{M}_{\triangle}(C,M)$ (see an example in Figure 12). Similarly as above, we proceed for even length cycle and in this case, we get $S\mathcal{M}_{\square}(C,M)$ (see an example in Figure 13). Therefore, when length of C is odd then the quadrangles can not repeat and when length of C is even then the triangles can not repeat. So, the triangles and quadrangles can not repeat simultaneously. Therefore, by combining above all the cases, the geometric carrier $|S|$ is $S(C,M),S\mathcal{M}_{\triangle}(C,M)$ or $S\mathcal{M}_{\square}(C,M)$. $\square$



Figure 10 : $S(C,M)$: Cylinder



Figure 12 : $S\mathcal{M}_{\triangle}(C,M)$



Figure 11



Figure 13 : $S\mathcal{M}_{\square}(C,M)$

By Lemma 2.14, the map M contains a $S(C,M),S\mathcal{M}_{\triangle}(C,M)$ or a $S\mathcal{M}_{\square}(C,M)$. For each of $S(C,M),S\mathcal{M}_{\triangle}(C,M)$ and $S\mathcal{M}_{\square}(C,M)$, we proceed with the argument as in Section 2.1 and hence, we get that the map M has either a $S(M)$ or a $S\mathcal{M}(M)$ representation.

We first assume that $S(M)$ denotes a representation of M. Then, $\partial S(M)=\{C,C'\}$ where C,C' are two cycles of type A_1. We recall definition of (r,s,k)-representation from [7] which is defined on the torus. The similar definition is also introduced in Section 2.1. Let $v\in V(C)$. We recall the following two paths in M as follows. Let $P(\ldots,v_{i-1},v_i,v_{i+1},\ldots)$ be a path in edge graph of M. The path P is of type A_2 [7] at vertex v_i if $lk(v_i)=C(\boldsymbol{a},v_{i-1},\boldsymbol{b},c,v_{i+1},d,e)$ implies $lk(v_{i+1})=C(\boldsymbol{a}_0,v_{i+2},\boldsymbol{b}_0,d,v_i,c,p)$ and $lk(v_i)=C(\boldsymbol{x},v_{i+1},\boldsymbol{z},l,v_{i-1},k,m)$ implies $lk(v_{i+1})=C(\boldsymbol{l},v_i,\boldsymbol{m},x,v_{i+2},g,z)$ for all i. Similarly, let $P(\ldots,w_{i-1},w_i,w_{i+1},\ldots)$ be a path in edge graph of M. The path P is of type A_3 at vertex w_i [7] if $lk(w_i)=C(\boldsymbol{a},w_{i-1},\boldsymbol{b},c,d,w_{i+1},e)$ implies $lk(w_{i+1})=C(\boldsymbol{a}_1,w_{i+2},\boldsymbol{b}_1,p,e,w_i,d)$ and $lk(w_i)=C(\boldsymbol{a}_2,w_{i+1},\boldsymbol{b}_2,p,e,w_{i-1},d)$ implies $lk(w_{i+1})=C(\boldsymbol{p},\ w_i,\boldsymbol{d},a_2,z_1,w_{i+2},b_2)$ for all i. We have two paths through v of types A_2 and A_3 (by definition). Let $L_1=C,L_2=P(a_1=v,\ldots,w),L_3=P(a_1=v,\ldots,w')$ denote two paths of types A_2 and A_3 respectively through v where $w,w'\in V(C')$. Let $C:=C(a_1,a_2,\ldots,a_r)$. Similarly as in Section 2.1, we say that C is a horizontal base cycle and a path is vertical if it is not homologous to C. Let faces incident with C be quadrangles and faces incident with C' be triangles in $S(M)$. Then, we make an another cut along L_3. Let s denote the number of cycles which are homologous to C along the path L_3. Let length$(L_3)=m$. Here, one can observe thjat the s is equal to the length of the path L_3, that is, $s=m$. Now length$(C)=r$ and number of horizontal cycles along L_3 is s. Thus, we get a planar polyhedral representation of $S(M)$ and it is denoted by (r,s)-*representation*. In the representation, we get identification of vertical sides in the natural manner but the identification of the horizontal sides needs some shifting with twist so that a vertex in the lower(base) side is identified with a vertex in the upper side. Let $C(a_{k+1}(=w_m), a_k, a_{k-1}, \ldots, a_k)$ denote the upper horizontal cycle in (r, s)-representation. So, the vertex a_{k+1} is the starting vertex of the upper horizontal cycle. Denote $k = \text{length}(P(a_{k+1}(=w_m), a_k, a_{k-1}, \ldots, a_k))$ in C. Hence, we represent the (r, s)-representation by (r, s, k)-*representation* and denote it by $K(r, s, k)$.

Assume that $S\mathcal{M}(M)$ denotes a representation of M. We cut along the boundaries of the Möbius strips as in Section 2.1. Hence, we get three components. These are a cylinder and two Möbius strips. We denote the cylinder by $S'(M)$ and two Möbius strips by $\mathcal{M}', \mathcal{M}''$ where $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}_{\triangle}(C, M), \mathcal{M}_{\Box}(L, M) \mid \text{ for some cycles } C \text{ and } L \text{ of type } A_1\}$. Thus, in this case, we denote the representation $S\mathcal{M}(M)$ by $K(S'(M), \mathcal{M}', \mathcal{M}'')$. Let l denote the length of a cycle of type A_1 in $S'(M)$. Then, the cycles $\partial\mathcal{M}', \partial\mathcal{M}''$ have same length l as $\partial S'(M) = \{\partial\mathcal{M}', \partial\mathcal{M}''\}$. Therefore, we get a $K(l, S'(M), \mathcal{M}', \mathcal{M}'')$ representation of $K(S'(M), \mathcal{M}', \mathcal{M}'')$. We combine above all arguments and hence, we have the following result.

Lemma 2.15. *The map M has a $K(r, s, k)$ or a $K(l, S'(M), \mathcal{M}', \mathcal{M}'')$ representation.*

We show in Lemma 2.16 that the M does not have both the representations. That is, we have

Lemma 2.16. *Let M_1 and M_2 be two maps on same number of vertices. Let $K_1(r, s, k)$ denote M_1 and $K_2(S'(M_2), \mathcal{M}'_2, \mathcal{M}''_2)$ denote M_2. Then, $M_1 \ncong M_2$.*

Proof. The map M_2 contains a Möbius strip $\mathcal{M}'_2$ which consists of either triangles or quadrangles. Let $C = \partial\mathcal{M}'_2$. Since M_1 does not contain Möbius strip which is bounded by any cycle of type A_1, so, there is no isomorphism map which maps C to a cycle of type A_1 in M_1. Therefore, $M_1 \ncong M_2$. ☐

From definition of A_1, we have one cycle of type A_1 through each vertex in M. It is discussed in [8] that the cycles of type A_1 in M have same length. Thus we have

Proposition 2.1. *([8]) The cycles of type A_1 in M have unique length.*

We study $K(r, s, k)$ and $K(S(M), \mathcal{M}', \mathcal{M}'')$, and classify them separately as they are non-isomorphic.

Classification of $K(r, s, k)$ on n vertices : We define admissible relations among r, s, k of $K(r, s, k)$ such that $K(r, s, k)$ represents a map after identifying its boundaries. The proof of the bellow results similarly as in Section 2.1. So, we have

Lemma 2.17. *The maps of type $\{3^3, 4^2\}$ of the form $K(r, s, k)$ exist if and only if the following holds : (i) $rs \geq 12$, (ii) $r \geq 3$, (iii) $s \geq 4$, and (iv) $k \in \{t : 0 \leq t \leq r-1\}$ if $s \geq 4$.*

Lemma 2.18. *Let $K(r_i, s_i, k_i)$ denote a (r_i, s_i, k_i)-representation of M_i on same number of vertices for $i \in \{1, 2\}$. Then, $M_1 \cong M_2$ if $(r_1, s_1, l_1) = (r_2, s_2, l_2)$ where $l_i \in \{k_i \ mod(2), (k_i + r_i) \ mod(2)\}$.*

Proof. The proof repeats similar argument as in Lemma 2.7. Let $r_1 = r_2, s_1 = s_2, k_1 = k_2$. Define an isomorphism map $f_1 : V(K(r_1, s_1, k_1)) \to V(K(r_2, s_2, k_2))$. So, $M_1 \cong M_2$ by f_1. Let $k_1 \neq k_2$. Repeat similar argument as in Lemma 2.7. Hence, $K(r_i, s_i, k_i)$ has either a $K(r_i, s_i, 0)$ or a $K(r_i, s_i, 1)$ representation. Hence, by $f_1, M_1 \cong M_2$. This completes the proof. ☐

Corollary 2.4. *Let $K(r_i,s_i,k_i)$ denote a (r_i,s_i,k_i)-representation of M_i on same number of vertices for $i \in \{1,2\}$. Then, $K(r_1,s_1,k_1) \ncong K(r_2,s_2,k_2)$ $\forall$ $r_1 \neq r_2, K(r_1,s_1,k_1) \ncong K(r_2,s_2,k_2)$ $\forall$ $s_1 \neq s_2, K(r_i,s_i,0) \cong K(r_i,s_i,k_i)$ for $2 \mid k_i$ or $2 \mid (k_i+r_i)$, and $K(r_i,s_i,1) \cong K(r_i,s_i,k_i)$ for $2 \mid (k_i-1)$ or $2 \mid (k_i+r_i-1)$.*

Classification of $K(l,t)$ on n vertices : Let $S\mathcal{M}(M)$ denote M on n vertices. By Lemma 2.15, the $S\mathcal{M}(M)$ has a $K(l,S'(M),\mathcal{M}',\mathcal{M}'')$ representation. Let $C' = \partial\mathcal{M}', C'' = \partial\mathcal{M}''$. Again, similarly as in the Proposition 2.1, length(C') = length(C''). Let $C' = \partial\mathcal{M}', C'' = \partial\mathcal{M}''$ such that $\partial S'(M) = \{C',C''\}$ for some cycles C',C'' of type A_1. Hence, we denote $K(l,S'(M),\mathcal{M}',\mathcal{M}'')$ by $K(l,t)$ if $l = length(C') = length(C'')$ and t denote the number of cycles of type A_1 in $S'(M)$.

Lemma 2.19. *The maps of the type $\{3^3,4^2\}$ of the form $K(l,t)$ exist if and only if the following holds : (i) $tl \geq 10$, (ii) $t \geq 2$ and $2 \mid t$, (iii) $l \geq 5$ if $t = 2$, (iv) $l \geq 4$ if $t \geq 4$, (v) $\mathcal{M}'$ and $\mathcal{M}''$ consists of only triangles if l is odd, (vi) $\mathcal{M}'$ and $\mathcal{M}''$ consists of only quadrangles if l is even.*

Proof. Let $K(l,S'(M),\mathcal{M}',\mathcal{M}'')$ denote a representation of M. It has a decompositions into a cylinder $S'(M)$ and two Möbius strips $\mathcal{M}'$ and $\mathcal{M}''$. By the similar argument as in Lemma 2.14, the cylinder $S'(M)$ contains two disjoint boundaries. So, $t \geq 2$. We repeat similar argument as in Lemma 2.8 and hence, we get $n = tl, 2 \mid t, l \mid n$ and $t \mid n$. Similarly we repeat similar arguments as in Lemma 2.17 for Case 3 and 4. By Proposition 2.1, the cycles of type A_1 have same length. Observe that by definition, when the length of the cycle is odd then by Lemma 2.14, $\mathcal{M}'$ and $\mathcal{M}''$ consist of only triangles. When the length of the cycle is even, then by Lemma 2.14, $\mathcal{M}'$ and $\mathcal{M}''$ consist of only quadrangles. This completes the proof. ☐

Let M_1 and M_2 be two maps on same number of vertices. Let $K(l_i,t_i)$ denote a representation of M_i. Then, we have

Lemma 2.20. *The $M_1 \cong M_2$ if $l_1 = l_2$.*

Proof. By Proposition 2.1, the M_i contains cycles of type A_1 of same length. We denote it by l. As in Lemma 2.18, let $n = sl$. So, the number of cycles of type A_1 in M_i is s. Let $C_1, \ldots, C_s$ denote the cycles of type A_1 in M_1. Let $C_1 = \partial\mathcal{M}_1'$ and $C_s = \partial\mathcal{M}_1''$. Again, let $L_1,\ldots,L_s$ denote the cycles of type A_1 in M_2. Let $L_1 = \partial\mathcal{M}_2'$ and $L_s = \partial\mathcal{M}_2''$. Since $l_1 = l_2$, so, the $\mathcal{M}_1', \mathcal{M}_1'', \mathcal{M}_2'$ and $\mathcal{M}_2''$ consist of either triangles or quadrangles. Therefore, we define an isomorphism map $f_1 : V(M_1) \to V(M_2)$ by $f_1(V(C_i)) = V(L_i)$ for $1 \leq i \leq s$ as in Lemma 2.18. Thus, $M_1 \cong M_2$. This completes the proof. ☐

Thus, we have

Corollary 2.5. *Let $K(r_i,s_i,k_i)$ denote a (r_i,s_i,k_i)-representation of M_i on same number of vertices for $i \in \{1,2\}$. Then, $K(l_1,t_1) \ncong K(l_2,t_2)$ $\forall$ $l_1 \neq l_2, K(l_1,t_1) \ncong K(l_2,t_2)$ $\forall$ $t_1 \neq t_2, M_i$ has an unique representation $K(l_i,t_i)$ for $2 \mid l_i$ and $n = l_it_i$, and M_i has an unique representation $K(l_i,t_i)$ for $2 \mid (l_i+1)$ and $n = l_it_i$.*

2.5 Maps of type $\{3^2,4,3,4\}$: Let M be a map of type $\{3^2,4,3,4\}$ on the Klein bottle. Let $P(\ldots,u_{i-1},u_i,u_{i+1},\ldots)$ be a path in edge graph of M. The path P is of type B_1 [7] if $lk(u_i) = C(\boldsymbol{a},u_{i+1},b,c,\boldsymbol{d},u_{i-1},e)$ implies $lk(u_{i-1}) = C(\boldsymbol{f},g,e,u_i,\boldsymbol{c},d,u_{i-2})$ and $lk(u_{i+1}) = C(\boldsymbol{e},a,k,u_{i+2},\boldsymbol{l},b,u_i)$, and $lk(u_i) = C(\boldsymbol{e},h,k,u_{i+1},\boldsymbol{l},b,u_{i-1})$ implies $lk(u_{i-1}) =$

$C(\boldsymbol{h}, u_i, b, c, \boldsymbol{d}, u_{i-2}, e)$ and $lk(u_{i+1}) = C(\boldsymbol{s}, u_{i+2}, t, l, \boldsymbol{b}, u_i, k)$ at u_i for all i. Let W be a maximal walk of type B_1 in M. Similarly as in Section 2.1, it is either a cycle or a closed walk. Observe that by definition, we have two paths of type B_1 through each vertex in M. Therefore, by the similar argument as in Section 2.2, the map M contains a cycle of type B_1 thorough each vertex of M. Let C be a cycle of type B_1 in M. Let S denote a set of faces which are incident with C. Then, we have

Lemma 2.21. *The* $|S|$ *is* $S(C, M)$ *or* $\mathcal{M}(C, M)$.

Proof. We first assume that the faces incident with C do not repeat. We repeat the similar argument as we have done in above preceding sections. Let $lk(w_1) = C(x_1, x_2, w_2, \boldsymbol{v_2}, v_1, w_r, \boldsymbol{x_r})$. We consider geometric carrier $|S|$ and cut along the path $P(x_1, w_1, v_1)$ of type B_1. Thus we have the following cases. Let path $P(x_1, w_1, v_1)$ identify with $P(x_1, w_1, v_1)$ with out any twist. In this case, we get a cylinder (for an example see in Figure 14). We denote it by $S(C, M)$. Let $P(x_1, w_1, v_1)$ identify with $P(x_1, w_1, v_1)$ with a twist. In this case, we get a Möbius strip (for an example see in Figure 15). We denote it by $\mathcal{M}(C, M)$.

Assume that the faces incident with C repeat. Let S' denote the set of faces incident with C and lie on one side of C. The set S' consists of both triangles and quadrangles. Similarly as in Lemma 2.2, each face of S' repeats exactly twice in the sequence of incident faces of C which are lie one side and belong in S'. So, let $F_1, F_2, \dots, F_k, F_1, F_2, \dots, F_k$ denote a sequence of faces which are incident with C for some k and belong to S'. That is $\{F_1, F_2, \dots, F_k, F_1, F_2, \dots, F_k\} = \{F_1, F_2, \dots, F_k\}$. Thus we have the following two cases. Let the cycle $C = C(w_1, \dots, w_{2r})$ be of even length. In this case, we consider link of w_{r+2} and hence, we get a face sequence either $\{3, 4, 3, 4\}$ or $\{3^2, 4, 3^2, 4\}$ (for example see in Figure 14 and 15). These are different from $\{3^2, 4, 3, 4\}$. So, length(C) is not even. Assume that the length of C is odd. In this case, the face sequence incident with C is either $\triangle'_1, \triangle'_1, \square_2, \dots, \square_{r-1}, \triangle'_r, \triangle''_r$ or $\square_1, \triangle'_2, \triangle''_2, \dots, \triangle'_{r-1}, \triangle''_{r-1}, \square_r$ ($\triangle', \square$ represent triangles and quadrangles respectively). In $\triangle'_1, \triangle''_1, \square_2, \dots, \square_{\frac{r-1}{2}}, \triangle'_{\frac{r+1}{2}}, \triangle''_{\frac{r+1}{2}}, \square_{\frac{r+3}{2}}, \dots, \square_{r-1}, \triangle'_r, \triangle''_r$ implies $\triangle'_1 = \triangle''_{\frac{r+1}{2}}$ and $\triangle''_1 = \square_{\frac{r+3}{2}}$ as $(\triangle'_1, \triangle'_1, \square_2, \dots, \square_{\frac{r-1}{2}}, \triangle'_{\frac{r+1}{2}}) = (\triangle''_{\frac{r+1}{2}}, \square_{\frac{r+3}{2}}, \dots, \square_{r-1}, \triangle'_r, \triangle''_r)$. But $\triangle''_1 \neq \square_{\frac{r+3}{2}}$. So, this case is not possible. Similarly as above, the sequence $\square_1, \triangle'_2, \triangle''_2, \dots, \triangle'_{r-1}, \triangle''_{r-1}, \square_r$ does not exist. Therefore, $|S| = S(C, M)$ (for example see in Figure 14) or $|S| = \mathcal{M}(C, M)$ (for example see in Figure 15). $\square$

So, by Lemma 2.21, M contains either of a cylinder $S(C, M)$ and a Möbius strip $\mathcal{M}(C, M)$. If $S(C, M) \subset M$ or $\mathcal{M}(C, M) \subset M$ then by the similar argument as in Section 2.1, the map M has either a $S(M)$ or a $S\mathcal{M}(M)$ representation. Let $S(M)$ denote M such that $\partial S(M) = \{C, C'\}$ for some cycles C, C' of type B_1. Let $v \in V(C)$. By definition, let $L = P(v, \dots, w)$ denote the path of type B_1 through v where $w \in V(C')$. We repeat the similar argument as in Section 2.4 which is done to define (r, s, k)-representation of M. Then, we take second cut along L where the starting adjacent face to the base horizontal cycle is a 4-gon. Thus, we get a representation of M and it is denoted by $K(r, s, k)$.

Again, let $S\mathcal{M}(M)$ denote M. Similarly as in the above preceding sections, we cut along the boundaries of the Möbius strips and hence, we get three components namely a cylinder $S'(M)$ and two Möbius strips $\mathcal{M}', \mathcal{M}''$ where $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}(L, M) \mid$ for some cycle L of type B_1 in $S\mathcal{M}(M)\}$. We denote this representation by $K(S'(M), \mathcal{M}', \mathcal{M}'')$. By combining above all cases, the map M has either a $K(r, s, k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation. We show in next lemma that every map M has a $K(r, s, k)$ representation.

Lemma 2.22. *The map* M *has a* $K(r, s, k)$ *representation.*

Proof. As above the map M has either a $K(r, s, k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation. Assume that $K(S'(M), \mathcal{M}', \mathcal{M}'')$ denotes M. We repeat similar process as in Lemma 2.10. We cut $\mathcal{M}(L, M)$ along a path P of type B_1, and hence, we get a planar polyhedral representation. Let denoted it by $\square(\mathcal{M})$. Similarly, we cut $S'(M)$ along two paths of type B_1. Hence, we get two planar polyhedral representations $\square'$ and $\square''$. And also, we cut $\mathcal{M}(L', M)$ along a path type B_1. Hence, we get an another polyhedral representation $\square(\mathcal{M}')$. In this process, we consider paths which are not homologous to the boundary cycles of the cylinder and Möbius strips through some fixed vertices. Now, similarly as in Lemma 2.10, we consider $\square(\mathcal{M})$ and $\square'$, and identify these along their common boundaries. Then, we consider $\square(\mathcal{M})$ and $\square''$, and identify these along their common boundaries. Thereafter, we consider $\square''$ and $\square(\mathcal{M}')$, and identify these along their common boundaries. Thus, we get a planar polyhedral representation $K(r', s', k')$ from $K(S'(M), \mathcal{M}', \mathcal{M}'')$ for some r', s', k'. Therefore, every map M has a $K(r, s, k)$ representation for some r, s, k. $\square$


x_1 x_2 x_3 x_4 x_{r-2} x_{r-1} x_r x_1
w_1 w_2 w_3 w_4 w_{r-2} w_{r-1} w_r w_1
v_1 v_2 v_3 v_4 v_{r-2} v_{r-1} v_r v_1

Figure 14 : $S(C, M)$: Cylinder


x_1 x_2 x_3 x_4 x_r x_{r+1} x_{r+2} x_{r+3} x_{2r-2} x_{2r-1} x_{2r} x_1
w_{r+3}
w_1 w_2 w_3 w_4 w_r w_{r+1} w_{r+2} w_{2r-2} w_{2r-1} w_{2r} w_1
w_{r+1} w_{r+2} w_{r+3} w_{r+4} w_2 w_1 w_2 w_3 w_{r-2} w_{r-1} w_r w_{r+1}

Figure 16


x_1 x_2 x_3 x_4 x_{r-2} x_{r-1} v_1
w_1 w_2 w_3 w_4 w_{r-2} w_{r-1} w_1
v_1 v_2 v_3 v_4 v_{r-2} v_{r-1} x_1

Figure 15


w_{r+4}
w_{r+1} w_{r+2} w_{r+3} w_{2r} w_1 w_2 w_3 w_{r-2} w_{r-1} w_r w_{r+1}
w_{r+3}
w_1 w_2 w_3 w_4 w_r w_{r+1} w_{r+2} w_{2r-2} w_{2r-1} w_{2r} w_1
v_1 v_2 v_3 v_4 v_r v_{r+1} v_{r+2} v_{r+3} v_{2r-2} v_{2r-1} v_{2r} v_1

Figure 17

As above, by definition, we have two paths through each vertex in M. This implies that we have two cycles of type B_1 upto homologous in M. Let C' and C'' denote two non-homologous cycles of type B_1 in M. As in Section 2.2, let C' homologous to the generator of $\mathbb{Z}$ and C_2 homologous to generator of $\mathbb{Z}_2$ of the fundamental group of M. As these are two different groups, so, there is no isomorphism which maps one cycle to another cycle. So, we consider only one cycle in the proof of Lemma 2.24. By the above Lemma 2.22, every map has a $K(r, s, k)$ representation. So, we study only $K(r, s, k)$ representations in bellow. Next we define admissible relations among r, s, k of $K(r, s, k)$ such that $K(r, s, k)$ represents a map after identifying its boundaries. The proof of next lemmas repeats similar argument as in Lemmas 2.6 and 2.18. Thus, we have

Lemma 2.23. *The maps of type* $\{3^2, 4, 3, 4\}$ *of the form* $K(r, s, k)$ *exist if and only if the following holds : (i)* $rs \geq 12$, *(ii)* $r \geq 4$, *(iii)* $2 \mid r$, *(iv)* $s \geq 3$, *(v)* $2 \nmid s$, *and (v)* $k \in \{2t \ : \ 0 \leq t \leq (\frac{r}{2} - 1)\}$ *if* $s \geq 3$.

Lemma 2.24. *Let* $K(r_i, s_i, k_i)$ *denote a* (r_i, s_i, k_i)*-representation of* M_i *on same number of vertices for* $i \in \{1, 2\}$. *Then,* $M_1 \cong M_2$ *if* $(r_1, s_1, l_1) = (r_2, s_2, l_2)$ *where* $l_i \in \{k_i \ mod(2), (k_i + r_i) \ mod(2)\}$.

Corollary 2.6. *Let* $K(r_i, s_i, k_i)$ *denote a* (r_i, s_i, k_i)*-representation of* M_i *on same number of vertices for* $i \in \{1, 2\}$. *Then,* $K(r_1, s_1, k_1) \not\cong K(r_2, s_2, k_2) \ \forall \ r_1 \neq r_2, K(r_1, s_1, k_1) \not\cong K(r_2, s_2, k_2) \ \forall \ s_1 \neq s_2, K(r_1, s_1, 0) \cong K(r_1, s_1, k_1)$ *for* $2 \mid k_1$ *or* $2 \mid (k_1 + r_1)$, *and* $K(r_1, s_1, 1) \cong K(r_1, s_1, k_1)$ *for* $2 \mid (k_1 - 1)$ *or* $2 \mid (k_1 + r_1 - 1)$.

2.6 Maps of type $\{4, 8^2\}$ **:** Let M be a map of type $\{4, 8^2\}$ on the Klein bottle. Through each vertex in M, we have a path of type Z_1 [7] in M. Let P be a maximal walk of type Z_1. By the similar argument as in Section 2.1, it is a cycle since degree is three. Let C be a cycle of type Z_1 in M. Let S denote a set of faces incident with C. Then, we have

Lemma 2.25. *The* $|S|$ *is* $S(C, M), \mathcal{M}(C, M)$ *or* $\mathcal{M}_{4,8}(C, M)$.

Proof. We repeat similar arguments as in Lemmas 2.14 and 2.21. By the similar argument as in Lemma 2.14, $|S|$ is either a cylinder $S(C, M)$ or a Möbius strip $\mathcal{M}_{4,8}(C, M)$. In this case, the $\mathcal{M}_{4,8}(C, M)$ denotes a Möbius strip which consists of 4-gons and 8-gons (the similar notion is also introduced in Section 2.1). Again, by the similar argument as in Lemma 2.21, $|S| = \mathcal{M}(C, M)$. Therefore, $|S| = S(C, M)$ or $|S| = \mathcal{M}(C, M)$ or $|S| = \mathcal{M}_{4,8}(C, M)$. This completes the proof. $\square$

So, by Lemma 2.25, $S(C, M) \subset M, \mathcal{M}(C, M) \subset M$ or $\mathcal{M}_{4,8}(C, M) \subset M$. We consider each of $S(C, M), \mathcal{M}(C, M)$ and $\mathcal{M}_{4,8}(C, M)$ and repeat the similar process as in Section 2.4, 2.5. Hence, we get either a $K(r, s, k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation of M. Next we show that the map M has a $K(r, s, k)$ representation.

Lemma 2.26. *The* M *has a* $K(r, s, k)$ *representation.*

Proof. Let $K(S'(M), \mathcal{M}', \mathcal{M}'')$ denote a representation of M. Then, $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}(C_1, M), \mathcal{M}_{4,8}(C_2, M) \mid$ for some cycles C_1, C_2 of type $Z_1\}$. We repeat the similar construction as in Lemma 2.22. Hence, we get a planar polyhedral representation $K(r, s, k)$ for some r, s, k. $\square$

Let C' and C'' be two non-homologous cycles in M. Similarly as in Section 2.5, there is no isomorphic maps which maps cycle C' to C''. Therefore, we study only $K(r, s, k)$ representations. Now we define admissible relations among r, s, k of $K(r, s, k)$ such that $K(r, s, k)$ represents a map after identifying its boundaries. The proof of next lemmas repeats similar argument as in Lemmas 2.6 and 2.18.

Lemma 2.27. *The maps of type* $\{4, 8^2\}$ *of the form* $K(r, s, k)$ *exist if and only if the following holds : (i)* $rs \geq 24$, *(ii)* $4 \mid r$, *(iii)* $s \geq 3$, *(iv)* $r \geq 8$, *(v)* $k \in \{4t+3 \ : \ 0 \leq t \leq \frac{r-4}{4}\}$ *if* $s \geq 3$ *and odd (vi)* $k \in \{(4t+2)(mod\ r) \ : \ 0 \leq t \leq \frac{r-4}{4}\}$ *if* $s \geq 4$ *and even.*

Lemma 2.28. *Let* $K(r_i, s_i, k_i)$ *denote a* (r_i, s_i, k_i) *-representation of* M_i *on same number of vertices for* $i \in \{1, 2\}$. *Then,* $M_1 \cong M_2$ *if* $(r_1, s_1, l_1) = (r_2, s_2, l_2)$ *where* $l_i \in \{k_i\ mod(2), (k_i + r_i)\ mod(2)\}$.

Corollary 2.7. *Let* $K(r_i, s_i, k_i)$ *denote a* (r_i, s_i, k_i)*-representation of* M_i *on same number of vertices for* $i \in \{1, 2\}$. *Then,* $K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2) \forall r_1 \neq r_2$, $K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2) \forall s_1 \neq s_2, K(r_1, s_1, 0) \cong K(r_1, s_1, k_1)$ *for* $2 \mid k_1$ *or* $2 \mid (k_1 + r_1), K(r_1, s_1, 1) \cong K(r_1, s_1, k_1)$ *for* $2 \mid (k_1 - 1)$ *or* $2 \mid (k_1 + r_1 - 1)$.

2.7 Maps of type $\{3, 6, 3, 6\}$ **:** Let M be a map of type $\{3, 6, 3, 6\}$ on the Klein bottle. Let W be a maximal walk of type X_1 [7] through each vertex in M. The maximal walk W is either a cycle or a closed walk as in Section 2.2. Similarly as in Section 2.1, the M contains a cycle of type X_1. Let C be a cycle of type X_1 in M. Let S denote a set of faces incident with C. Then, we have

Lemma 2.29. *The* $|S|$ *is* $S(C, M), \mathcal{M}(C, M)$ *or* $\mathcal{M}_{3,6}(C, M)$.

Proof. We argue similarly as in Lemmas 2.14 and 2.21. Assume that faces which are incident with C are all distinct. In this case, by the similar argument as in Lemmas 2.14 and 2.21, $|S| = S(C, M)$ and $|S| = \mathcal{M}(C, M)$ respectively. Assume that faces which are incident with C are not all distinct. Let $\triangle_1, F_2, \triangle_3, F_4, \ldots, F_{i-1}, \triangle_i, F_{i+1}, \ldots \triangle_{k-1}, F_k$ denote a sequence of faces which are incident with C in ordered and lie on one side of C where F_i denote a 6-gon for each i. Since $\triangle_i \cap F_{i-1}, \triangle_i \cap F_{i+1}$ and $\triangle_i \cap C$ are each an edge, so, the triangles are distinct in the sequence. But the faces F_is' are repeating in the sequence. Since each 6-gon F_i has exactly one edge remaining to identify, so, by similar argument as in Lemma 2.21, F_i repeats exactly once in the sequence. So, the geometric carrier $|S| = \mathcal{M}_{3,6}(C, M)$ and which consists of 3-gons and 6-gons. Therefore, the $|S|$ is either of $S(C, M), \mathcal{M}(C, M)$ and $\mathcal{M}_{3,6}(C, M)$. This completes the proof. □

So, by Lemma 2.29, $S(C, M) \subset M$ or $\mathcal{M}(C, M) \subset M$ or $\mathcal{M}_{3,6}(C, M) \subset M$ for some cycle C of type X_1 in M. We repeat similar argument as in Sections 2.1 and 2.5, the map M has either a $S(M)$ representation which is bounded by two identical cycles with opposite direction or a $S\mathcal{M}(M)$ which is bounded by two Möbius strips $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}(C, M), \mathcal{M}_{3,6}(L, M)\}$ from each of $S(C, M), \mathcal{M}(C, M)$ and $\mathcal{M}_{3,6}(C, M)$. Let $S(M)$ denote a representation of M such that $\partial S(M) = \{C, C'\}, \triangle \in F(M)$ and $C \cap \triangle$ is an edge. By definition of X_1, we have three paths of type X_1 through three edges of $\triangle$. Let L_1, L_2, L_3 be three paths of type X_1. We repeat the similar argument as in Section 2.5 to define a (r, s, k)-representation of a map M. Let $L_1 = C$. We take a cut along L_3. Since length(C) = length(C') and $C = C'$, hence, we get a planar polyhedral representation which is bounded by L_3, C, L_3 and C'. Since $C = C'$, so, it defines a $K(r, s, k)$ representation. Let $S\mathcal{M}(M)$ denote a representation of M. We cut along the boundaries of the Möbius strips. We get three components. These are namely a cylinder $S'(M)$ and two Möbius strips $\mathcal{M}', \mathcal{M}''$ where $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}(C, M), \mathcal{M}_{3,6}(C, M)\}$ for some cycle C of type X_1. We denote this representation by $K(S'(M), \mathcal{M}', \mathcal{M}'')$. By combining above all cases, the map M has either a $K(r, s, k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation and these are non-isomorphic. We study both the representations separately in bellow.

Classification of $K(r, s, k)$ on n vertices : Let $\triangle$ in M and $e_1, e_2, e_3 \subset \triangle$. By definition of X_1, we have three paths and similarly as in Section 2.1, let C be a cycle of type X_1 such that $e_1 \subset C$ and W be a closed walk of type X_1 such that that $e_2, e_3 \subset C$. So, there is one cycle of type X_1 upto homologous. We define admissible relations among r, s, k of $K(r, s, k)$ such that $K(r, s, k)$ represents a map after identifying its boundaries. The proof of next lemmas repeats similar argument as in Lemmas 2.6 and 2.18.

Lemma 2.30. *The maps of type $\{3, 6, 3, 6\}$ of the form $K(r, s, k)$ on n vertices exist if and only if the following holds : (i) $\frac{3}{2}rs \geq 27$, (ii) $s \geq 3$, (iii) $2 \mid r$, (iv) $r \geq 6$, and (v) $k \in \{2t + 1 \ : \ 0 \leq t \leq \frac{r}{2} - 1\}$.*

Lemma 2.31. *Let $K(r_i, s_i, k_i)$ denote a (r_i, s_i, k_i) -representation of M_i on same number of vertices for $i \in \{1, 2\}$. Then, $M_1 \cong M_2$ if $(r_1, s_1, l_1) = (r_2, s_2, l_2)$ where $l_i \in \{k_i \ mod(2), (k_i + r_i) \ mod(2)\}$.*

Corollary 2.8. *Let $K(r_i, s_i, k_i)$ denote a (r_i, s_i, k_i)-representation of M_i on same number of vertices for $i \in \{1, 2\}$. Then, $K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2) \forall r_1 \neq r_2, K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2) \forall s_1 \neq s_2, K(r_1, s_1, 0) \cong K(r_1, s_1, k_1)$ for $2 \mid k_1$ or $2 \mid (k_1 + r_1)$, and $K(r_1, s_1, 1) \cong K(r_1, s_1, k_1)$ for $2 \mid (k_1 - 1)$ or $2 \mid (k_1 + r_1 - 1)$.*

Classification of $K(l,t)$ on n vertices : Let $S\mathcal{M}(M)$ denote M. Then, by the above argument, $K(S'(M),\mathcal{M}',\mathcal{M}'')$ has $S\mathcal{M}(M)$ representation. Let $C'=\partial\mathcal{M}', C''=\partial\mathcal{M}''$. Then, length$(C')$ = length(C'') as C' and C'' are boundaries of $S'(M)$. Let l = length(C'). Now $\mathcal{M}',\mathcal{M}'' \in \{\mathcal{M}(L',M),\mathcal{M}_{3,6}(L'',M) \mid$ for some cycles L' and L'' of type $X_1\}$. Let $\mathcal{M}'=\mathcal{M}(L',M)$ and $\mathcal{M}''=\mathcal{M}_{3,6}(L'',M)$. By definition and Lemma 2.29, $C''=L''$ and length(C') = 2×length(L'). Thus, the map M contains cycles of type X_1 of two different lengths. In this case, we denote $K(S'(M),\mathcal{M}',\mathcal{M}'')$ by $K(l_1,l_2,S'(M),\mathcal{M}',\mathcal{M}'')$ where $l_2=l$ and $2l_1=l_2$. When $\mathcal{M}'=\mathcal{M}_{3,6}(L',M)$ and $\mathcal{M}''=\mathcal{M}_{3,6}(L'',M)$ then we denote $K(S'(M),\mathcal{M}',\mathcal{M}'')$ by $K(l,S'(M),\mathcal{M}',\mathcal{M}'')$. The above all notions are used in the next lemma.

Lemma 2.32. *The maps of type $\{3,6,3,6\}$ of the form $K(\frac{l}{2},l,S'(M),\mathcal{M}',\mathcal{M}'')$ exist if and only if the following holds :*

(1) $t\geq 2, n=\frac{3}{2}tl+\frac{l}{2}$ where $4\mid l$ and $l\geq 12$ if $\mathcal{M}'=\mathcal{M}_{3,6}(C',M)$ and $\mathcal{M}''=\mathcal{M}_{3,6}(C'',M)$.

(2) $t\geq 1, n=l(t+2)$ where $2\mid l$ and $l\geq 10$ if $\mathcal{M}'=\mathcal{M}(L',M)$ and $\mathcal{M}''=\mathcal{M}(L'',M)$ for some cycles L' and L'' of type X_1.

(3) $t\geq 1, n=l(t+\frac{5}{4})$ where $4\mid l$ and $l\geq 12$ if $\mathcal{M}'=\mathcal{M}_{3,6}(C',M)$ and $\mathcal{M}''=\mathcal{M}(L'',M)$.

Proof. The proof repeats similar argument as in Section 2.1. In all the cases, if we consider velues of t,l not in the given range then we get some vertex whose link is not a cycle. □

Lemma 2.33. *Let $K(S'(M_i),\mathcal{M}_i',\mathcal{M}_i'')$ denote M_i on same number of vertices. Then, we have the following.*

(1) Let $K(l_i,S'(M_i),\mathcal{M}_i',\mathcal{M}_i'')$ denote M_i. Let $\mathcal{M}_i'=\mathcal{M}_{3,6}(C_i',M_i)$ and $\mathcal{M}_i''=\mathcal{M}_{3,6}(C_i'',M_i)$. Then, $M_1\cong M_2$ if $l_1=l_2$.

(2) Let $K(l_i,\frac{l_i}{2},S'(M_i),\mathcal{M}_i',\mathcal{M}_i'')$ denote M_i. Let $\mathcal{M}_i'=\mathcal{M}(C_i',M_i)$ and $\mathcal{M}_i''=\mathcal{M}(C_i'',M_i)$. Then, $M_1\cong M_2$ if $l_1=l_2$.

(3) Let $\mathcal{M}_i'=\mathcal{M}_{3,6}(C_i',M_i)$ and $\mathcal{M}_i''=\mathcal{M}(C_i'',M_i)$. Then, $M_1\cong M_2$ if $l_1=l_2$.

Proof. Let $K(l_i,S'(M_i),\mathcal{M}_i',\mathcal{M}_i'')$ denote M_i and $\mathcal{M}_i'=\mathcal{M}_{3,6}(C',M)$ and $\mathcal{M}_i''=\mathcal{M}_{3,6}(C'',M)$. By Lemma 2.32, $\frac{3}{2}t_1l_1+\frac{1}{2}l_1=n=\frac{3}{2}t_2l_2+\frac{1}{2}l_2$ as $l_1=l_2$. Hence, $t_1=t_2=s$ (say). Let $C_1,\ldots,C_s$ denote the cycles of type X_1 in M_1. Let $C_1=\partial\mathcal{M}_1'$ and $C_s=\partial\mathcal{M}_1''$. Again, let $L_1,\ldots,L_s$ denote the cycles of type X_1 in M_2. Let $C_i=C(u(i,1),\ldots,u(i,l_1))$ and $L_i=C(v(i,1),\ldots,v(i,l_2))$. Let $L_1=\partial\mathcal{M}_2'$ and $L_s=\partial\mathcal{M}_2''$. We repeat similar argument as in Lemma 2.7. Hence, we define an isomorphism map $f_1: V(M_1)\to V(M_2)$ by $f_1(V(C_i))=V(L_i)$ and $f_1(u(j,1))=v(j,1)$ for $1\leq j\leq l_1$ and $1\leq i\leq s$. So, $M_1\cong M_2$. Similar argument we repeat for other two cases. If two maps have two different representation in the above three cases, then, they are non-isomorphic as they have non-isomorphic Möbius strips. This completes the proof. □

2.8 Maps of type $\{3,12^2\}$: Let M be a map of type $\{3,12^2\}$ on the Klein bottle. Let W be a maximal walk of type G_1 [7] in edge graph of M. Then, it is either a cycle or a closed walk (see similar argument as in Section 2.7). Similarly as in Section 2.7, the map M contains a cycle of type G_1. Let C be a cycle of type G_1 in M. Let S denote a set of faces which are incident with C. Then, $|S|=S(C,M)$ or $|S|=\mathcal{M}(C,M)$ or $|S|=\mathcal{M}_{3,12}(C,M)$ (see similar argument as in Lemma 2.29). Hence, either $S(C,M)\subset M,\mathcal{M}(C,M)\subset M$ or a $\mathcal{M}_{3,12}(C,M)\subset M$. Consider each of $S(C,M),\mathcal{M}(C,M)$ and $\mathcal{M}_{3,12}(C,M)$ and we proceed similarly as in Section 2.7, hence, we get either a $K(r,s,k)$ or a $K(S'(M),\mathcal{M}',\mathcal{M}'')$

representation of M and these are non-isomorphic. We study both the representations separately in bellow.

Classification of $K(r, s, k)$ on n vertices : Let $\triangle$ in $F(M)$. Similarly as in the previous Section 2.7, let C be a cycle of type G_1 which contains an edge of $\triangle$ and W be a closed walk of type G_1 which contains other two edges of $\triangle$. Hence, there is one cycle of type G_1 upto homologous in M. We define admissible relations among r, s, k of $K(r, s, k)$ in the next lemma and its proof follows from the similar arguments as in Lemma 2.6. Then, the next isomorphism lemma whose proves follows from similar argument as in Lemma 2.18.

Lemma 2.34. *The maps of type $\{3, 12^2\}$ of the form $K(r, s, k)$ on n vertices exist if and only if the following holds : (i) $n = \frac{3}{2}rs$, (ii) $4 \mid r$, (iii) $6 \mid n$, (iv) $s \geq 3$, (v) $r \geq 12$, and (vi) $k_1 \in \{4t + 2 \ : \ 0 \leq t \leq \frac{r-4}{4}\}$.*

Lemma 2.35. *Let $K(r_i, s_i, k_i)$ denote a (r_i, s_i, k_i)-representation of M_i on same number of vertices for $i \in \{1, 2\}$. Then, $M_1 \cong M_2$ if $(r_1, s_1, l_1) = (r_2, s_2, l_2)$ where $l_i \in \{k_i \ mod(2), (k_i + r_i) \ mod(2)\}$.*

Corollary 2.9. *$K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2)$ if $r_1 \neq r_2, K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2)$ if $s_1 \neq s_2, K(r_1, s_1, 0) \cong K(r_1, s_1, k_1)$ if $2 \mid k_1$ or $2 \mid (k_1 + r_1)$, and $K(r_1, s_1, 1) \cong K(r_1, s_1, k_1)$ if $2 \mid (k_1 - 1)$ or $2 \mid (k_1 + r_1 - 1)$.*

Classification of $K(l, t)$ on n vertices : Let M be a map of type $\{3, 12^2\}$ on n vertices. We repeat similar arguments and notions of Section 2.7 in bellow. Now we have the next lemma and isomorphism lemma. The proof of the lemma follows similar argument as in Lemma 2.32 and the proof of the isomorphism lemma follows from similar argument as in Lemma 2.33.

Lemma 2.36. *The maps of type $\{3, 12^2\}$ of the form $K(\frac{l}{2}, l, S'(M), \mathcal{M}', \mathcal{M}'')$ exist if and only if the following holds :*

(1) $t \geq 2, n = \frac{3}{2}tl + \frac{l}{2}$ where $8 \mid l$ and $l \geq 24$ if $\mathcal{M}' = \mathcal{M}_{3,12}(C', M)$ and $\mathcal{M}'' = \mathcal{M}_{3,12}(C'', M)$.

(2) $t \geq 1, n = l(t + 2)$ where $4 \mid l$ and $l \geq 20$ if $\mathcal{M}' = \mathcal{M}(L', M)$ and $\mathcal{M}'' = \mathcal{M}(L'', M)$ for some cycles L' and L'' of types G_1.

(3) $t \geq 1, n = l(t + \frac{5}{4})$ where $4 \mid l$ and $l \geq 24$ if $\mathcal{M}' = \mathcal{M}_{3,12}(C', M)$ and $\mathcal{M}'' = \mathcal{M}(L'', M)$.

Lemma 2.37. *Let $K(S'(M_i), \mathcal{M}'_i, \mathcal{M}''_i)$ represent M_i on n vertices. Then, we have the following cases :*

(1) Let $K(l_i, S'(M_i), \mathcal{M}'_i, \mathcal{M}''_i)$ represents M_i such that $\mathcal{M}'_i = \mathcal{M}_{3,12}(C'_i, M_i)$ and $\mathcal{M}''_i = \mathcal{M}_{3,12}(C''_i, M_i)$. Then $M_1 \cong M_2$ if $l_1 = l_2$.

(2) Let $K(l_i, \frac{l_i}{2}, S'(M_i), \mathcal{M}'_i, \mathcal{M}''_i)$ represents M_i such that $\mathcal{M}'_i = \mathcal{M}(C'_i, M_i)$ and $\mathcal{M}''_i = \mathcal{M}(C''_i, M_i)$. Then $M_1 \cong M_2$ if $l_1 = l_2$.

(3) Let $\mathcal{M}'_i = \mathcal{M}_{3,12}(C'_i, M_i)$ and $\mathcal{M}''_i = \mathcal{M}(C''_i, M_i)$. Then $M_1 \cong M_2$ if $l_1 = l_2$.

2.9 Maps of type $\{3^4, 6\}$: Let M be a map of type $\{3^4, 6\}$ on the Klein bottle. Let P be a maximal walk of type Y_1 [7]. It is either a cycle or a closed walk as in Section 2.1. As in Section 2.7, the map M contains a cycle, say C of type Y_1 in M. Let S denote a set of faces incident with C. Then, similarly as in Lemma 2.29, $|S| = S(C, M)$. Next we show that there is no map of type $\{3^4, 6\}$ on the Klein bottle.

Theorem 2.1. *There is no map of type $\{3^4, 6\}$ on the Klein bottle.*

Proof. By the above argument, let $S(C, M) \subset M$ for some cycle C of type Y_1. We consider $S(C, M)$ and repeat with the arguments as in Sections 2.4, 2.5. In this process, let $S(M)$ denote M.

Claim : The cylindrical representation $S(M)$ does not exist in M.

Let $\partial S(M) = \{C', C''\}$ such that $C' = C''$ and C' identifies with C'' with a twist. Let $C' = C(w_1, \ldots, w_r)$. By definition of Y_1, C' divides number of triangles at w_i one of $0 : 4, 1 : 3$ and $2 : 2$ ratios (here, $a : b$ ratio is to mean a number of triangles lie on one side and b number of triangles lie on the other side of the cycle at a vertex). Let $P(w_{j-1}, w_j, w_{j+1}) \subset C', C''$ for some j as $C' = C''$. Let $i \in \{0, 1, 2\}$. If $P \subset C'$ then P divides incident triangles one of $i : 4-i, 1+i : 3-i$ and $2+i : 2-i$ ratios for some fixed i. If $P \subset C''$ then P divides the incident triangles one of $i : 4-i, 2+i : 2-i$ or $1+i : 3-i$ ratios for some fixed i. Now we have the following cases. Observe that by definition of Y_1, if ratio $i : 4-i$ is at $w_i \in C'$ then $i : 4-i$ is at $w_i \in C''$ in $S(M)$. Again, if ratio $1+i : 3-i$ is at $w_i \in C'$ then $2+i : 2-i$ is at $w_i \in C''$. Similarly again, if ratio $2+i : 2-i$ is at $w_i \in C'$ then $1+i : 3-i$ is at $w_i \in C''$. In later two cases, it is impossible to identify boundaries of $S(M)$ at w_i as the $lk(w_i)$ divides incident faces into two different ratios. So, the C' can not identify with C'' with a twist. So, $S(M)$ does not exist. Therefore, there is no map of type $\{3^4, 6\}$ on the Klein bottle. This completes the proof. $\square$

2.10 Maps of type $\{4, 6, 12\}$ **:** Let M be a map of type $\{4, 6, 12\}$ on the Klein bottle. Let P be a maximal walk of type H_1 [7] in M. It is either a cycle or a closed walk (see the similar argument as in Section 2.7). Since degree of each vertex is three, it follows that the map M has a cycle of type H_1 as in Section 2.7. Let C be a cycle of type H_1 in M. Then, the geometric carrier of the faces which are incident with C is either $S(C, M)$ or $\mathcal{M}_{4,12}(C, M)$ (see the similar argument as in Lemma 2.29). Thus, either $S(C, M) \subset M$ or $S\mathcal{M}_{4,12}(C, M) \subset M$. We consider each of $S(C, M)$ and $S\mathcal{M}_{4,12}(C, M)$, and repeat the similar arguments as in Section 2.5. Hence, we get either a $K(r, s, k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation of M.

Classification of $K(r, s, k)$ **on** n **vertices :** We define admissible relations among r, s, k of $K(r, s, k)$ such that $K(r, s, k)$ represents a map after identifying its boundaries. The proof of next lemma repeats similar argument as in Lemma 2.6 and the proof of Lemma 2.39 follows from similarly as in Lemma 2.18. Thus, we have

Lemma 2.38. *The maps of tyep* $\{4, 6, 12\}$ *of the form* $K(r, s, k)$ *on* n *vertices exist if and only if the following holds : (i)* $n = rs \geq 48$, *(ii)* $2 \mid s$, *(iii)* $6 \mid r$, *(iv)* $s \geq 4$, *(v)* $12 \mid n$, *(vi)* $r \geq 12$, *and (vii)* $k \in \{6t + 4 \ : \ 0 \leq t \leq \frac{r-6}{6}\}$.

Lemma 2.39. *Let* $K(r_i, s_i, k_i)$ *denote a* (r_i, s_i, k_i)*-representation of* M_i *on same number of vertices for* $i \in \{1, 2\}$. *Then,* $M_1 \cong M_2$ *if* $(r_1, s_1, l_1) = (r_2, s_2, l_2)$ *where* $l_i \in \{k_i \ mod(2), (k_i + r_i) \ mod(2)\}$.

Corollary 2.10. *The* $K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2)$ *if* $r_1 \neq r_2, K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2)$ *if* $s_1 \neq s_2, K(r_1, s_1, 0) \cong K(r_1, s_1, k_1)$ *if* $2 \mid k_1$ *or* $2 \mid (k_1 + r_1)$, *and* $K(r_1, s_1, 1) \cong K(r_1, s_1, k_1)$ *if* $2 \mid (k_1 - 1)$ *or* $2 \mid (k_1 + r_1 - 1)$.

Classification of $K(l, t)$ **on** n **vertices :** Let M be a map of type $\{4, 6, 12\}$ on n vertices. Let $S\mathcal{M}(M)$ denote M. Then, it has a $K(l, S'(M), \mathcal{M}', \mathcal{M}'')$ representation (see similar argument as in Section 2.4). Let C' and C'' denote two boundary cycles of $\mathcal{M}', \mathcal{M}''$ respectively. Then, $C' = \partial\mathcal{M}', C'' = \partial\mathcal{M}''$ and length(C') = length(C''). Let

$C' = \partial\mathcal{M}', C'' = \partial\mathcal{M}''$ and $\{C_1, C_2\} = \partial S'(M)$. Let $l = length(C') = length(C'')$ and t denote the number of cycles in $S'(M)$ which are of type H_1. The proof of next lemmas repeats similar argument as in Lemmas 2.19 and 2.33.

Lemma 2.40. *The maps of type* $\{4, 6, 12\}$ *of the form* $K(S'(M), \mathcal{M}', \mathcal{M}'')$ *on* n *vertices exist if and only if the following holds : (i)* $t \geq 2$, *(ii)* $2 \mid t$, *(iii)* $n = tl$ *where* $12 \mid l, l \geq 24$ *if* $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}_{4,12}(C, M)\}$.

Let $K(l_i, S'(M_i), \mathcal{M}'_i, \mathcal{M}''_i)$ denote a representation of M_i on n vertices where $\mathcal{M}'_i, \mathcal{M}''_i \in \{\mathcal{M}_{4,12}(C, M_i) \mid$ for some cycle C of type $H_1\}$. Then, we have :

Lemma 2.41. *The map* $M_1 \cong M_2$ *if* $l_1 = l_2$.

Let t_i denote the number of cycles of type H_1 in M_i. Then $K(l_i, t_i)$ denotes $K(S'(M_i), \mathcal{M}'_i, \mathcal{M}''_i)$ for $i \in \{1, 2\}$. Then, we have

Corollary 2.11. *The* $K(l_1, t_1) \ncong K(l_2, t_2)$ *if* $l_1 \neq l_2$ *and* $K(l_1, t_1) \ncong K(l_2, t_2)$ *if* $t_1 \neq t_2$.

2.11 Maps of type $\{3, 4, 6, 4\}$ **:** Let M be a map of type $\{3, 4, 6, 4\}$ on the Klein bottle. It has a walk of type W_1 [7] in M through each vertex in M. Similarly as in Section 2.1, let C be a cycle of type W_1 in M. Let S denote a set of faces which are incident with C. Then $|S|$ is one of $S(C, M), \mathcal{M}_{3,4}(C, M)$ and $\mathcal{M}_{4,6}(C, M)$ as in Lemma 2.29. So, the map M contains either $S(C, M), \mathcal{M}_{3,4}(C, M)$ or $\mathcal{M}_{4,6}(C, M)$. We consider each of $S(C, M), \mathcal{M}_{3,4}(C, M)$ and $\mathcal{M}_{4,6}(C, M)$, and repeat the similar arguments as in Sections 2.4, 2.5. Hence, we get either a $K(r, s, k)$ or a $K(S'(M), \mathcal{M}', \mathcal{M}'')$ representation of M. We study both the representations in bellow.

Classification of $K(r, s, k)$ **on** n **vertices :** We define admissible relations among r, s, k of $K(r, s, k)$ such that $K(r, s, k)$ represents a map after identifying its boundaries. It's proof repeats similar argument as in Lemma 2.6 and the proof of next Lemma 2.43 follows from similar argument as in Lemma 2.18. Thus, we have

Lemma 2.42. *The maps of type* $\{3, 4, 6, 4\}$ *of the form* $K(r, s, k)$ *on* n *vertices exist if and only if the following holds : (i)* $n = rs \geq 24$, *(ii)* $3 \mid r$, *(iii)* $2 \mid s$, *(iv)* $s \geq 4$, *(v)* $r \geq 6$, *and (vi)* $k \in \{3t + 2 \ : \ 0 \leq t \leq \frac{r-3}{3}\}$.

Lemma 2.43. *Let* $K(r_i, s_i, k_i)$ *denote a* (r_i, s_i, k_i)*-representation of* M_i *on same number of vertices for* $i \in \{1, 2\}$. *Then,* $M_1 \cong M_2$ *if* $(r_1, s_1, l_1) = (r_2, s_2, l_2)$ *where* $l_i \in \{k_i\ mod(2), (k_i + r_i)\ mod(2)\}$.

Corollary 2.12. $K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2)$ *if* $r_1 \neq r_2$, $K(r_1, s_1, k_1) \ncong K(r_2, s_2, k_2)$ *if* $s_1 \neq s_2, K(r_1, s_1, 0) \cong K(r_1, s_1, k_1)$ *if* $2 \mid k_1$ *or* $2 \mid (k_1 + r_1)$, *and* $K(r_1, s_1, 1) \cong K(r_1, s_1, k_1)$ *if* $2 \mid (k_1 - 1)$ *or* $2 \mid (k_1 + r_1 - 1)$.

Classification of $K(l, t)$ **on** n **vertices :** We repeat the similar argument as in Section 2.10. Let $K(l, S'(M), \mathcal{M}', \mathcal{M}'')$ denote a representation of M. Let C' and C'' denote two boundary cycles of $\mathcal{M}', \mathcal{M}''$ respectively. Let $C' = \partial\mathcal{M}', C'' = \partial\mathcal{M}''$ and $\partial S'(M) = \{C', C''\}$. Let $l = length(C') = length(C'')$ and t denote the number of cycles of type W_1 in $S'(M)$. Let $\mathcal{M}', \mathcal{M}'' \in \{\mathcal{M}_{3,4}(L', M), \mathcal{M}_{4,6}(L'', M)\}$ for some cycles L' and L'' of type W_1. Then we have the following lemmas.

Lemma 2.44. *The maps of type* $\{3, 4, 6, 4\}$ *of the form* $K(S'(M), \mathcal{M}', \mathcal{M}'')$ *on* n *vertices exist if and only if the following holds :*

(1) $t \geq 2, n = tl$ *where* $6 \mid (l - 3)$ *and* $l \geq 9$ *if* $\mathcal{M}' = \mathcal{M}_{3,4}(C', M)$ *and* $\mathcal{M}'' = \mathcal{M}_{3,4}(C'', M)$*.*

(2) $t \geq 2, n = lt$ *where* $t \geq 2, 6 \mid l$ *and* $l \geq 12$ *if* $\mathcal{M}' = \mathcal{M}_{4,6}(L', M)$ *and* $\mathcal{M}'' = \mathcal{M}_{4,6}(L'', M)$*.*

(3) $K(S'(M), \mathcal{M}', \mathcal{M}'')$ *not exists if* $\mathcal{M}' = \mathcal{M}_{3,4}(C', M)$ *and* $\mathcal{M}'' = \mathcal{M}_{4,6}(L', M)$*.*

Proof. We repeat similar argument as in Lemma 2.19 for first two cases. If there exists a map M which has a $K(l, S'(M), \mathcal{M}', \mathcal{M}'')$ representation such that $\mathcal{M}' = \mathcal{M}_{3,4}(C', M)$ and $\mathcal{M}'' = \mathcal{M}_{4,6}(L', M)$. Then by the above cases, $6 \mid (l - 3)$ and $6 \mid l$. But there is no such l which satisfies both the conditions. So, M does not exist. This completes the proof. □

The proof of Lemma 2.45 repeats similar argument as in Lemma 2.33. Thus, we have

Lemma 2.45. *Let* $K(l_i, S'(M_i), \mathcal{M}'_i, \mathcal{M}''_i)$ *denote* M_i *on* n *vertices. Then, we have the following :*

(1) Let $\mathcal{M}'_i = \mathcal{M}_{3,4}(C'_i, M_i)$ *and* $\mathcal{M}''_i = \mathcal{M}_{3,4}(C''_i, M_i)$*. Then* $M_1 \cong M_2$ *if* $l_1 = l_2$*.*

(2) Let $\mathcal{M}'_i = \mathcal{M}_{4,6}(C'_i, M_i)$ *and* $\mathcal{M}''_i = \mathcal{M}_{4,6}(C'_i, M_i)$*. Then* $M_1 \cong M_2$ *if* $l_1 = l_2$*.*

3 Proofs

Proof of Theorem 1.1 : From Section 2.1, maps of type $\{3^6\}$ have either $K(r, s, k)$ or $K(l, t)$ representation for some r, s, k, l, t. By Lemma 2.6, it follows that $rs \geq 9, r \geq 3, s \geq 3$. Let $m = s, r = \frac{n}{m}$. Then $m = s \geq 3, r = \frac{n}{m} \geq 3$ i.e. $n \geq 3m$. Now we have following cases. If $gcd(r, 2) = 1$ i.e. $gcd(n, 2m) = m$ then $r + k$ is even for odd $k \in \{t : 0 \leq t \leq r - 1\}$. That is, k and $r + k$ are both even for $gcd(r, 2) = 1$ and $k \in \{t : 0 \leq t \leq r - 1\}$. In this case, we get only one map which is isomorphic to $K(r, s, 0)$. If $gcd(r, 2) = 2$ i.e. $gcd(n, 2m) = 2m$ then $r + k, k$ are odd for odd $k \in \{t : 0 \leq t \leq r - 1\}$. In this case, we get two maps where one is isomorphic to $K(r, s, 1)$ and another is isomorphic to $K(r, s, 0)$. Therefore, in this case, we get two maps. By Lemma 2.8, $K(l, t)$ exists if (i) $tl \geq 10$, (ii) $t \geq 2$, (iii) $2 \nmid l$ and $l \geq 5$. Let $m = t$. Then $m \geq 2, m \mid n, n \geq 5m$. If $gcd(n, 2m) = m$ i.e. $\frac{n}{m}$ is even then it has an unique representation $K(\frac{n}{m}, m)$. If $gcd(n, 2m) = 2m$ then there is no map by Lemma 2.8. Since $K(r, s, k)$ and $K(l, t)$ are non-isomorphic by Lemma 2.4, so, by combining above all three cases, if the number of maps, say $i(n)_{\{3^6\}}$ on n vertices then $i(n)_{\{3^6\}} = \sum_{i=1,2} i \times |\{(m, n) \mid m \geq 3, n \geq 3m,\ gcd(n, 2m) = i \times m\}| + |\{(m, n) \mid m \geq 2, m \mid n, n \geq 5m,\ gcd(n, 2m) = m\}|$.

From Section 2.2, by Lemma 2.11, rs ≥ 9, r ≥ 3, s ≥ 3 and $k \in \{t : 0 \leq t \leq r - 1\}$ for $s \geq 3$. Let $m = s, r = \frac{n}{m}$. Similarly as above, if $k \in \{t : 0 \leq t \leq r - 1\}$ then k and $r + k$ both are even for $gcd(r, 2) = 1$. In this case, we get only one map which is isomorphic to $K(r, s, 0)$. If $gcd(r, 2) = 2$ then similarly, $r + k, k$ are odd for odd $k \in \{t : 0 \leq t \leq r - 1\}$. In this case, we get two maps where one is isomorphic to $K(r, s, 1)$ and another is isomorphic to $K(r, s, 0)$. Thus, we get two non-isomorphic maps. By combining above all two cases, if number of maps $i(n)_{\{4^4\}}$ (say) on n vertices then $i(n)_{\{4^4\}} = \sum_{i=1,2} i \times |\{(m, n) \mid m \geq 3, n \geq 3m,\ gcd(n, 2m) = i \times m\}|$.

Let $i(2n)_{\{6^3\}}$ denote the number of maps of type $\{6^3\}$ on $2n \geq 14$ vertices. Then, by duality, $i(2n)_{\{6^3\}} = i(n)_{\{3^6\}}$ as number of faces in map of type $\{3^6\}$ is equal to the number of vertices in map of type $\{6^3\}$.

Similarly as above by Lemmas 2.17, 2.19, let $i(n)_{\{3^3,4^2\}}$ denote the number of maps of type $\{3^3,4^2\}$ on n vertices. Then $i(n)_{\{3^3,4^2\}} = \sum_{i=1,2} i \times |\{(m,n) \mid m \geq 4, 2 \mid m, n \geq 3m,\ gcd(n,2m) = i \times m\}| + |\{(m,n) \mid m \geq 5, n = 2m\}| + |\{(m,n) \mid m \geq 4, n \geq 4m, 2m \mid n\}|$.

By Lemma 2.23, let $i(n)_{\{3^2,4,3,4\}}$ denote the number of maps of type $\{3^2,4,3,4\}$ on n vertices. Then $i(n)_{\{3^2,4,3,4\}} = |\{(m,n) \mid m \geq 3, 2 \nmid m, 2m \mid n, n \geq 12\}|$.

By Lemma 2.27, let $i(n)_{\{4,8^2\}}$ denote the number of maps of type $\{4,8^2\}$ on n vertices. Then $i(n)_{\{4,8^2\}} = |\{(l,m,n) \mid m \geq 3, 2 \mid m, n \geq 8m, 4m \mid n, 0 \leq l \leq (\frac{n}{4m} - 1)\}| + |\{(l,m,n) \mid m \geq 3, 2 \nmid m, n \geq 8m, 4m \mid n, 0 \leq l \leq (\frac{n}{4m} - 1)\}|$.

By Lemmas 2.30, 2.32, let $i(n)_{\{3,6,3,6\}}$ denote the number of maps of type $\{3,6,3,6\}$ on n vertices. Then $i(n)_{\{3,6,3,6\}} = |\{(l,m,n) \mid m \geq 3, 3m \mid n, n \geq 9m, 0 \leq l \leq (\frac{n}{3m} - 1)\}| + |\{(m,n)\ | m \geq 2, 6m+2 \mid n, n \geq 6(3m+1)\}| + |\{(m,n) \mid m \geq 1, 2(m+2) \mid n, n \geq 10(m+2)\}| + |\{(m,n) \mid m \geq 1, 4m+5 \mid n, n \geq 12(4m+5)\}|$.

By Lemmas 2.34, 2.36, let $i(n)_{\{3,6,3,6\}}$ denote the number of maps of type $\{3,12^2\}$ on n vertices. Then $i(n)_{\{3,12^2\}} = |\{(l,m,n) \mid m \geq 3, 3m \mid n, n \geq 9m, 0 \leq l \leq (\frac{n}{6m} - 1)\}| + |\{(m,n) \mid m \geq 2, 4(3m+1) \mid n, n \geq 12(3m+1)\}| + |\{(m,n) \mid m \geq 1, 4(m+2) \mid n, n \geq 20(m+2)\}| + |\{(m,n) \mid m \geq 1, 4m+5 \mid n, n \geq 24(4m+5)\}|$.

By Theorem 2.1, there is no map of type $\{3^4,6\}$ on n vertices. So, $i(n)_{\{3^4,6\}} = 0$ for all n.

By Lemmas 2.38, 2.40, let $i(n)_{\{4,6,12\}}$ denote the number of maps of type $\{4,6,12\}$ on n vertices. Then $i(n)_{\{4,6,12\}} = \{(l,m,n) \mid m \geq 4, 2 \mid m, 6m \mid n, n \geq 12m, 0 \leq l \leq (\frac{n}{6m} - 1)\}| + |\{(m,n) \mid m \geq 2, 2 \mid m, 12m \mid n, n \geq 24m\}|$.

By Lemmas 2.42, 2.44, let $i(n)_{\{3,4,6,4\}}$ denote the number of maps of type $\{3,4,6,4\}$ on n vertices. Then $i(n)_{\{3,4,6,4\}} = |\{(l,m,n) \mid m \geq 4, 2 \mid m, 3m \mid n, n \geq 6m, 0 \leq l \leq (\frac{n}{3m} - 1)\}| + |\{(m,n) \mid m \geq 2, 6m \mid (n-3m), n \geq 9m\}| + |\{(m,n) \mid m \geq 2, 6m \mid n, n \geq 12m\}|$.
☐

Proof of Theorem 1.2 : The proof follows from the Sections 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 2.10, 2.11. Let M be a map of type $\{3^6\}$ on the Klein bottle with n vertices. By Lemmas 2.6, 2.8, we consider all admissible $K(r,s,k)$ and $K(l,t)$ representations of M on n vertices. Then, we classify them by Lemmas 2.7 and 2.9. Let $r_1, r_2 \mid n$. By Corollary 2.1, 2.2, $K(r_1,s,k) \ncong K(r_2,s,k), K(r_1,t) \ncong K(r_2,t)$ if $r_1 \neq r_2$ for all s,k,t, and $K(r,s,k)$ is isomorphic to either of $K(r,s,0), K(r,s,1)$. Hence, every map of type $\{3^6\}$ on n vertices is isomorphic to one of $K(r,s,0), K(r,s,1)$ and $K(r,s)$ for some $r \mid n, s = \frac{n}{r}$.

Similarly by Corollary 2.3, $K(r_1,s,k) \ncong K(r_2,s,k)$ if $r_1 \neq r_2$ for all s,k, and every map of type $\{4^4\}$ on n vertices is isomorphic to one of $K(r,s,0), K(r,s,1)$ for some $r \mid n$ and $s = \frac{n}{r}$.

By Corollary 2.4, 2.5, $K(r_1,s,k) \ncong K(r_2,s,k)$ and $K(r_1,t) \ncong K(r_2,t)$ if $r_1 \neq r_2$ for all s,k,t, and $K(r,s,k)$ is isomorphic to either of $K(r,s,0), K(r,s,1)$. So, every map of type $\{3^3,4^2\}$ on n vertices is isomorphic to one of $K(r,s,0), K(r,s,1), K(r,s)$ for some $r \mid n$ and $s = \frac{n}{r}$.

By Corollary 2.6, $K(r_1,s,k) \ncong K(r_2,s,k)$ if $r_1 \neq r_2$ for all s,k, and every map of type $\{3^2,4,3,4\}$ on n vertices is isomorphic to one of $K(r,s,0), K(r,s,1)$ for some $r \mid n$ and $s = \frac{n}{r}$.

By Corollary 2.7, $K(r_1,s,k) \ncong K(r_2,s,k)$ if $r_1 \neq r_2$ for all s,k, and every map of type $\{4,8^2\}$ on n vertices is isomorphic to one of $K(r,s,0), K(r,s,1)$ for some $r \mid n$ and $s = \frac{n}{r}$.

By Corollary 2.8 and Lemma 2.33, $K(r_1,s,k) \ncong K(r_2,s,k)$ and $K(r_1,t) \ncong K(r_2,t)$ if $r_1 \neq r_2$ for all s,k,t, and $K(r,s,k)$ is isomorphic to either of $K(r,s,0), K(r,s,1)$. So, every map of type $\{3,6,3,6\}$ on n vertices is isomorphic to one of $K(r,s,0), K(r,s,1)$ and $K(r,s)$

for some $r \mid n$ and $s \in \{\frac{2n}{3r}, \frac{2n}{3r+1}, \frac{4n}{4r+5}\}$.

By Corollary 2.9 and Lemma 2.37, $K(r_1, s, k) \ncong K(r_2, s, k)$ and $K(r_1, t) \ncong K(r_2, t)$ if $r_1 \neq r_2$ for all s, k, t, and $K(r, s, k)$ is isomorphic to either of $K(r, s, 0), K(r, s, 1)$. So, every map of type $\{3, 12^2\}$ on n vertices is isomorphic to one of $K(r, s, 0), K(r, s, 1)$ and $K(r, s)$ for some $r \mid n$ and $s \in \{\frac{2n}{3r}, \frac{2n}{3r+1}, \frac{4n}{4r+5}\}$.

By Corollary 2.10 and 2.11, $K(r_1, s, k) \ncong K(r_2, s, k)$ and $K(r_1, t) \ncong K(r_2, t)$ if $r_1 \neq r_2$ for all s, k, t, and $K(r, s, k)$ is isomorphic to either of $K(r, s, 0), K(r, s, 1)$. So, every map of type $\{4, 6, 12\}$ on n vertices is isomorphic to one of $K(r, s, 0), K(r, s, 1), K(r, s)$ for some $r \mid n$ and $s = \frac{n}{r}$.

By Corollary 2.12 and Lemma 2.45, $K(r_1, s, k) \ncong K(r_2, s, k)$ and $K(r_1, t) \ncong K(r_2, t)$ if $r_1 \neq r_2$ for all s, k, t, and $K(r, s, k)$ is isomorphic to either of $K(r, s, 0), K(r, s, 1)$. So, every map of type $\{3, 4, 6, 4\}$ on n vertices is isomorphic to one of $K(r, s, 0), K(r, s, 1), K(r, s)$ for some $r \mid n$ and $s = \frac{n}{r}$. $\square$

4 Acknowledgement

The first author is supported by post doctoral scholarship of Harish Chandra Research Institute, Department of Atomic Energy, Government of India. Work of second author is partially supported by SERB, DST grant No. SR/S4/MS:717/10.

References

[1] A. Altshuler, Hamiltonian Circuits in Some Maps on the Torus, Discrete Math. 4 (1972) 299-314.

[2] A. Altshuler, Construction and enumeration of regular maps on the torus, Discrete Math. 4 (1973) 201-217.

[3] J. A. Bondy and U. S. R. Murthy, Graph Theory, GTM 244, Springer, 2008.

[4] U. Brehm and W. Kühnel, Equivelar maps on the torus, European J. Combin. 29 (2008) 1843-1861.

[5] B. Datta and D. Maity, Semi-equivelar and vertex-transtive maps of torus. (preprint)

[6] W. Kurth, Enumeration of Platonic maps on the torus, Discrete Math. 61 (1986) 71-83.

[7] D. Maity and A. K. Upadhyay, Hamiltonian Cycle in Semi-Equivelar Maps on the Torus, (arXiv:1308.6717).

[8] D. Maity and A. K. Upadhyay, Enumeration of Semi-Equivelar Maps on the Torus, (arXiv:1311.0105).

[9] D. Maity and A. K. Upadhyay, On enumeration of a class of maps on Klein bottle, (arXiv:1509.04519)

[10] S. Negami, Uniqueness and faithfulness of embedding of toroidal graphs, Discrete Math. 44 (1983) 161-180.

[11] A. K. Tiwari and A. K. Upadhyay, An Enumeration of Semi-Equivelar Maps on Torus and Klein bottle, Math. Slovaca (to appear).